

LYAPUNOV EXPONENTS AND SPECTRAL ANALYSIS OF ERGODIC SCHRÖDINGER OPERATORS: A SURVEY OF KOTANI THEORY AND ITS APPLICATIONS

DAVID DAMANIK

Dedicated to Barry Simon on the occasion of his 60th birthday.

ABSTRACT. The absolutely continuous spectrum of an ergodic family of one-dimensional Schrödinger operators is completely determined by the Lyapunov exponent as shown by Ishii, Kotani and Pastur.

Moreover, the part of the theory developed by Kotani gives powerful tools for proving the absence of absolutely continuous spectrum, the presence of absolutely continuous spectrum, and even the presence of purely absolutely continuous spectrum.

We review these results and their recent applications to a number of problems: the absence of absolutely continuous spectrum for rough potentials, the absence of absolutely continuous spectrum for potentials defined by the doubling map on the circle, and the absence of singular spectrum for the subcritical almost Mathieu operator.

1. INTRODUCTION

Schrödinger operators with ergodic potentials have enjoyed quite some popularity for several decades now. This is in no small part due to Barry Simon's contributions to the field, through research articles on the one hand, but also through survey articles and his way of putting his personal stamp on results and conjectures on the other hand. Ergodic Schrödinger operators continue to be dear to him as seven of the fifteen Schrödinger operator problems he singles out in [60] for further investigation in the 21st century deal with them. Moreover, the immense activity in the area of ergodic Schrödinger operators is reflected by the fact that the ratio 7/15 improves to 3/4 when it comes to the problems from that list that have been solved so far. One may say that this is due to the uneven distribution of difficulty among these fifteen problems, but this is balanced by the fact that of the remaining four ergodic problems at least three are very hard and that further progress should be expected on some of the remaining non-ergodic problems.

In the area of ergodic Schrödinger operators there are several powerful methods (e.g., fractional moment) and analyses (multi-scale) but few theories (Kotani). What appears to be wordplay wants to express the fact that Kotani theory is distinguished from the other greats by its immensely general scope. It really is a theory that applies to the class of all ergodic operators and it is central in many ways. In

Date: May 18, 2006.

2000 Mathematics Subject Classification. Primary 82B44, 47B80; Secondary 47B36, 81Q10.

Key words and phrases. Ergodic Schrödinger Operators, Lyapunov Exponents.

The author was supported in part by NSF grant DMS-0500910.

addition, Kotani theory has played a crucial role in the solution of two of the four recently solved 21st century problems.

Our goal here is to present the core parts of Kotani theory with more or less complete proofs and to discuss several recent applications of the theory to a number of concrete classes of models for which, whenever possible, we at least outline the main ideas that go into the proofs of the results we mention.

Suppose (Ω, μ) is a probability measure space, $T : \Omega \rightarrow \Omega$ is an invertible ergodic transformation, and $f : \Omega \rightarrow \mathbb{R}$ is bounded and measurable. We define potentials,

$$V_\omega(n) = f(T^n \omega), \quad \omega \in \Omega, \quad n \in \mathbb{Z},$$

and the corresponding discrete Schrödinger operators in $\ell^2(\mathbb{Z})$,

$$(1) \quad [H_\omega \psi](n) = \psi(n+1) + \psi(n-1) + V_\omega(n)\psi(n).$$

We will call $\{H_\omega\}_{\omega \in \Omega}$ an ergodic family of Schrödinger operators.

Examples. (a) Quasi-periodic potentials: $\Omega = \mathbb{T}^d = \mathbb{R}^d/\mathbb{Z}^d$, μ is the normalized Lebesgue measure on \mathbb{T}^d , and $T\omega = \omega + \alpha$ is some ergodic shift (i.e., $1, \alpha_1, \dots, \alpha_d$ are rationally independent).¹

(b) Potentials defined by the skew shift: $\Omega = \mathbb{T}^2$, μ is the normalized Lebesgue measure on \mathbb{T}^2 , and $T(\omega_1, \omega_2) = (\omega_1 + \omega_2, \omega_2 + \alpha)$ for some irrational α .

(c) Potentials defined by the doubling map: $\Omega = \mathbb{T}$, μ is the normalized Lebesgue measure on \mathbb{T} , and $T\omega = 2\omega$.²

(d) Potentials defined by the left shift: $\Omega = I^\mathbb{Z}$, where I is a compact subset of \mathbb{R} , $\mu = \mathbb{P}^\mathbb{Z}$, where \mathbb{P} is a Borel probability measure on I , and $[T\omega](n) = \omega(n+1)$.

The following pair of results, proven in [50, 56], shows that for ergodic families of Schrödinger operators, the spectrum and the spectral type are deterministic in the sense that they are constant μ -almost surely.

Theorem 1 (Pastur 1980). *There exists a set $\Sigma \subset \mathbb{R}$ such that for μ -almost every ω , $\sigma(H_\omega) = \Sigma$.*

Theorem 2 (Kunz-Souillard 1980). *There are sets $\Sigma_{ac}, \Sigma_{sc}, \Sigma_{pp} \subset \mathbb{R}$ such that for μ -almost every ω , $\sigma_\bullet(H_\omega) = \Sigma_\bullet$, $\bullet \in \{ac, sc, pp\}$.*

Thus, in the spectral analysis of a given ergodic family of Schrödinger operators, a fundamental problem is the identification of the sets $\Sigma, \Sigma_{ac}, \Sigma_{sc}$, and Σ_{pp} .

The almost sure spectrum, Σ , is completely described by the integrated density of states as shown by Avron and Simon [9]. Denote the restriction of H_ω to $[0, N-1]$ with Dirichlet boundary conditions by $H_\omega^{(N)}$. For $\omega \in \Omega$ and $N \geq 1$, define measures $dk_{\omega, N}$ by placing uniformly distributed point masses at the eigenvalues $E_\omega^{(N)}(1) <$

¹What we call quasi-periodic here is more general than the notion of quasi-periodicity as defined in [7], for example, where a quasi-periodic potential is almost periodic with a finitely generated frequency module. In particular, a quasi-periodic potential as defined here is not necessarily almost periodic, that is, the translates of a given quasi-periodic potential are not necessarily precompact in $\ell^\infty(\mathbb{Z})$. We allow discontinuous sampling functions f here because we want to include Fibonacci-type potentials.

²Strictly speaking, this example does not fall within our general framework as T is not invertible, but potentials of this kind have been studied in several works and it is possible to tweak the model a little to fit it in the framework above.

$\dots < E_\omega^{(N)}(N)$ of $H_\omega^{(N)}$, that is,

$$\int f(E) dk_{\omega,N}(E) = \frac{1}{N} \sum_{n=1}^N f(E_\omega^{(N)}(n)).$$

Then, it can be shown that for μ -almost every $\omega \in \Omega$, the measures $dk_{\omega,N}$ converge weakly to a non-random measure dk , called the *density of states measure*, as $N \rightarrow \infty$. The function k defined by

$$k(E) = \int \chi_{(-\infty, E]}(E') dk(E')$$

is called the *integrated density of states*. It is not hard to show that

$$(2) \quad \int f(E) dk(E) = \mathbb{E}(\langle \delta_0, f(H_\omega) \delta_0 \rangle)$$

for bounded measurable f . Here, $\mathbb{E}(\cdot)$ denotes integration with respect to the measure μ , that is, $\mathbb{E}(g) = \int g(\omega) d\mu(\omega)$. Thus, the density of states measure is given by an average of the spectral measures associated with H_ω and δ_0 . The T -invariance of μ then implies the following result:

Theorem 3 (Avron-Simon 1983). *The almost sure spectrum is given by the points of increase of k , that is, $\Sigma = \text{supp}(dk)$.*

There is a similarly general description of the set Σ_{ac} in terms of the Lyapunov exponent. Let $E \in \mathbb{C}$ and

$$(3) \quad A^E(\omega) = \begin{pmatrix} E - f(\omega) & -1 \\ 1 & 0 \end{pmatrix}.$$

Define the *Lyapunov exponent* $\gamma(E)$ by

$$\gamma(E) = \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}(\log \|A_n^E(\omega)\|),$$

where $A_n^E(\omega) = A^E(T^{n-1}\omega) \cdots A^E(\omega)$.

The integrated density of states and the Lyapunov exponent are related by the Thouless formula (see, e.g., [23, Theorem 9.20]), which reads

$$(4) \quad \gamma(E) = \int \log |E - E'| dk(E').$$

The significance of the transfer matrices $A_n^E(\omega)$ is that a sequence u solves the difference equation

$$(5) \quad u(n+1) + u(n-1) + V_\omega(n)u(n) = Eu(n)$$

if and only if

$$\begin{pmatrix} u_n \\ u_{n-1} \end{pmatrix} = A_n^E(\omega) \begin{pmatrix} u_0 \\ u_{-1} \end{pmatrix}$$

for every n . Since $\det A^E(\omega) = 1$, we always have $\gamma(E) \geq 0$. Let us define

$$\mathcal{Z} = \{E : \gamma(E) = 0\}$$

By general principles, $\mathcal{Z} \subseteq \Sigma$.

For a subset S of \mathbb{R} , the essential closure of S is given by

$$\overline{S}^{\text{ess}} = \{E \in \mathbb{R} : \text{Leb}(S \cap (E - \varepsilon, E + \varepsilon)) > 0 \text{ for every } \varepsilon > 0\}$$

Then, the following theorem combines results from [37, 43, 56].³

Theorem 4 (Ishii 1973, Pastur 1980, Kotani 1984). *The almost sure absolutely continuous spectrum is given by the essential closure of the set of energies for which the Lyapunov exponent vanishes, that is, $\Sigma_{ac} = \overline{\mathcal{Z}}^{\text{ess}}$.*

While there is an analog of Theorem 3 for higher-dimensional ergodic Schrödinger operators, Theorem 4 is, by its very nature, a strictly one-dimensional result. It is one of the great challenges for researchers in the area of ergodic Schrödinger operators to develop effective tools for the study of the absolutely continuous spectrum in higher dimensions. That said, Theorem 4 holds in virtually all one-dimensional and quasi-one-dimensional situations: see Kotani [43] for continuous one-dimensional Schrödinger operators (see also Kirsch [41] for a useful extension), Minami [54] for generalized Sturm-Liouville operators, Kotani-Simon [49] for discrete and continuous Schrödinger operators with matrix-valued potentials, and Geronimo [31] and Geronimo-Teplyaev [32] for orthogonal polynomials on the unit circle.

2. THE DESCRIPTION OF THE AC SPECTRUM

In this section we discuss the main ideas that go into the proof of Theorem 4. The proof naturally breaks up into the proof of two inclusions.

The inclusion “ \subseteq ” was proved by Ishii and Pastur. The other inclusion was proved by Kotani and is a much deeper result. In fact, the proof of the Ishii-Pastur half of the result we give below is based on more modern techniques and shows that this half is really an immediate consequence of the general theory of one-dimensional Schrödinger operators.

There are at least three different proofs of the Ishii-Pastur half of Theorem 4 in the literature. One of them uses the existence of generalized eigenfunctions; compare Cycon et al. [23]. The second one, due to Deift and Simon [30], is close in spirit to, and uses techniques from, Kotani’s proof of the other half of the result. Finally, there are two somewhat related proofs given by Buschmann [18] and Last and Simon [51], which are both either directly or indirectly based on a result of Gilbert and Pearson that describes a support of the singular spectrum of a Schrödinger operator with fixed (non-random) potential in terms of subordinate solutions. We will follow the argument from Buschmann’s paper below.

We first recall a central result from Gilbert and Pearson’s subordinacy theory [33, 34]. Consider the discrete Schrödinger operator H in $\ell^2(\mathbb{Z})$ with potential V and solutions of the difference equation

$$(6) \quad u(n+1) + u(n-1) + V(n)u(n) = Eu(n).$$

A non-zero solution u of (6) is called *subordinate* at $\pm\infty$ if for every linearly independent solution \tilde{u} of (6), we have

$$\frac{\sum_{n=0}^{N-1} |u(\pm n)|^2}{\sum_{n=0}^{N-1} |\tilde{u}(\pm n)|^2} \rightarrow 0 \text{ as } N \rightarrow \infty.$$

Let

$$S = \{E \in \mathbb{R} : (6) \text{ has solutions } u_+ \text{ and } u_- \text{ such that } u_{\pm} \text{ is subordinate at } \pm\infty\}.$$

³To be more precise, the discrete version of the Kotani half of this result here can be found in the paper [59] by Simon and the work of Ishii and Pastur was preceded by closely related work by Casher and Lebowitz [19].

Then, S has zero weight with respect to the absolutely continuous part of any spectral measure, that is,

$$(7) \quad \mathcal{P}^{(\text{ac})}(S) = 0.$$

Proof of the Ishii-Pastur half of Theorem 4. Note that when $\gamma(E) > 0$, Oseledec' Theorem [55] says that for almost every ω , there are solutions $u_+(E, \omega)$ and $u_-(E, \omega)$ of (5) such that $u_\pm(E, \omega)$ is exponentially decaying, and hence subordinate, at $\pm\infty$. Applying Fubini's theorem, we see that for μ -almost every ω , the set of $E \in \mathbb{R} \setminus \mathcal{Z}$ for which the property just described fails, has zero Lebesgue measure. In other words, for these ω 's, $\mathbb{R} \setminus \mathcal{Z} \subseteq S_\omega$ up to a set of zero Lebesgue measure. Since sets of zero Lebesgue measure have zero weight with respect to the absolutely continuous part of any spectral measure, we obtain from (7) that for μ -almost every ω ,

$$\mathcal{P}_\omega^{(\text{ac})}(\mathbb{R} \setminus \mathcal{Z}) = 0.$$

This shows that for μ -almost every ω , $\sigma_{\text{ac}}(H_\omega) \subseteq \overline{\mathcal{Z}}^{\text{ess}}$. \square

Let us now turn to the Kotani half of Theorem 4. Kotani worked in the continuum setting. Carrying his results over to the discrete case is not entirely straightforward and it was worked out by Simon [59] whose proof we give below. Given $z \in \mathbb{C}_+ = \{z \in \mathbb{C} : \Im z > 0\}$ and $\omega \in \Omega$, there are (up to a multiplicative constant) unique solutions $u_\pm(n, \omega)$ of

$$(8) \quad u(n+1) + u(n-1) + V_\omega(n)u(n) = zu(n)$$

such that u_\pm is square-summable at $\pm\infty$. (Take $u_\pm(n, \omega) = \langle \delta_n, (H_\omega - z)^{-1} \delta_1 \rangle$ near $\pm\infty$ to show existence; uniqueness follows from constancy of the Wronskian.) Note that $u_\pm(0, \omega) \neq 0$ for otherwise z would be a non-real eigenvalue of a self-adjoint half-line operator. Thus, we can define

$$(9) \quad m_\pm(z, \omega) = -\frac{u_\pm(\pm 1, \omega)}{u_\pm(0, \omega)}.$$

Clearly,

$$(10) \quad m_\pm(z, T^n \omega) = -\frac{u_\pm(n \pm 1, \omega)}{u_\pm(n, \omega)}.$$

By Oseledec' Theorem, we have for μ -almost every ω ,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \left| \frac{u_\pm(n, \omega)}{u_\pm(0, \omega)} \right| = -\gamma(z).$$

By (10),

$$\log \left| \frac{u_\pm(n, \omega)}{u_\pm(0, \omega)} \right| = \sum_{m=0}^{n-1} \log |m_\pm(z, T^{\pm m} \omega)|$$

and hence Birkhoff's ergodic theorem implies

$$(11) \quad \mathbb{E}(\log |m_\pm(z, \omega)|) = -\gamma(z).$$

Proposition 2.1. *We have that*

$$\mathbb{E} \left(\log \left(1 + \frac{\Im z}{\Im m_\pm(z, \omega)} \right) \right) = 2\gamma(z).$$

Proof. By the difference equation (8) that u_{\pm} obeys,

$$(12) \quad m_{\pm}(z, T^n \omega) = V_{\omega}(n) - z - [m_{\pm}(z, T^{n \mp 1} \omega)]^{-1}.$$

Taking imaginary parts,

$$\Im m_{\pm}(z, \omega) = -\Im z - \Im ([m_{\pm}(z, T^{\mp 1} \omega)]^{-1}).$$

Dividing by $\Im m_{\pm}(z, \omega)$,

$$1 = -\frac{\Im z}{\Im m_{\pm}(z, \omega)} - \frac{\Im ([m_{\pm}(z, T^{\mp 1} \omega)]^{-1})}{\Im m_{\pm}(z, \omega)}.$$

Taking the logarithm,

$$\begin{aligned} \log \left(1 + \frac{\Im z}{\Im m_{\pm}(z, \omega)} \right) &= \log (-\Im ([m_{\pm}(z, T^{\mp 1} \omega)]^{-1})) - \log (\Im m_{\pm}(z, \omega)) \\ &= \log \left(\frac{\Im m_{\pm}(z, T^{\mp 1} \omega)}{|m_{\pm}(z, T^{\mp 1} \omega)|^2} \right) - \log (\Im m_{\pm}(z, \omega)) \end{aligned}$$

Taking expectations and using invariance,

$$\begin{aligned} \mathbb{E} \left(\log \left(1 + \frac{\Im z}{\Im m_{\pm}(z, \omega)} \right) \right) &= \mathbb{E} \left(\log \left(\frac{\Im m_{\pm}(z, T^{\mp 1} \omega)}{|m_{\pm}(z, T^{\mp 1} \omega)|^2} \right) - \log (\Im m_{\pm}(z, \omega)) \right) \\ &= -2 \mathbb{E} (\log |m_{\pm}(z, \omega)|) \\ &= 2\gamma(z), \end{aligned}$$

where we used (11) in the last step. \square

Denote

$$b(z, \omega) = m_+(z, \omega) + m_-(z, \omega) + z - V_{\omega}(0).$$

Proposition 2.2. *We have*

$$\mathbb{E} \left(\Im \left(\frac{1}{b(z, \omega)} \right) \right) = -\frac{\partial \gamma(z)}{\partial (\Im z)}.$$

Proof. It follows from (12) that

$$(13) \quad \frac{u_-(1, \omega)}{u_-(0, \omega)} = m_-(z, \omega) + z - V_{\omega}(0).$$

It is not hard to check that for $n \leq m$,

$$(14) \quad G_{\omega}(n, m; z) := \langle \delta_n, (H_{\omega} - z)^{-1} \delta_m \rangle = \frac{u_-(n, \omega) u_+(m, \omega)}{u_+(1, \omega) u_-(0, \omega) - u_-(1, \omega) u_+(0, \omega)}.$$

From (9), (13), (14), we get

$$(15) \quad -G_{\omega}(0, 0; z)^{-1} = m_+(z, \omega) + m_-(z, \omega) + z - V_{\omega}(0) = b(z, \omega).$$

The definition of $G_{\omega}(n, m; z)$ gives

$$(16) \quad \mathbb{E}(G_{\omega}(0, 0; z)) = \int \frac{1}{E' - z} dk(E').$$

Thus,

$$\begin{aligned}
\mathbb{E} \left(\Im \left(\frac{1}{b(z, \omega)} \right) \right) &= -\Im \mathbb{E} (G_\omega(0, 0; z)) \\
&= -\Im \int \frac{1}{E' - z} dk(E') \\
&= -\frac{\partial}{\partial(\Im z)} \int \log |z - E'| dk(E') \\
&= -\frac{\partial \gamma(z)}{\partial(\Im z)},
\end{aligned}$$

where we used (15), (16), and the Thouless formula (4). \square

Denote

$$n_\pm(z, \omega) = \Im m_\pm(z, \omega) + \frac{1}{2} \Im z.$$

Proposition 2.3. *We have that*

$$(17) \quad \mathbb{E} \left(\frac{1}{n_\pm(z, \omega)} \right) \leq \frac{2\gamma(z)}{\Im z}$$

and

$$(18) \quad \mathbb{E} \left(\frac{\left[\frac{1}{n_+} + \frac{1}{n_-} \right] \cdot [(n_+ - n_-)^2 + (\Re b)^2]}{|b|^2} \right) \leq 4 \left[\frac{\gamma(z)}{\Im z} - \frac{\partial \gamma(z)}{\partial(\Im z)} \right].$$

Proof. For $x \geq 0$, consider the function $A(x) = \log(1+x) - \frac{x}{1+\frac{x}{2}}$. Clearly, $A(0) = 0$ and $A'(x) = \frac{1}{1+x} - \frac{1}{1+x+\frac{x^2}{4}} \geq 0$. Therefore,

$$(19) \quad \log(1+x) \geq \frac{x}{1+\frac{x}{2}} \quad \text{for all } x \geq 0.$$

Thus,

$$\begin{aligned}
\mathbb{E} \left(\frac{1}{n_\pm(z, \omega)} \right) &= \mathbb{E} \left(\frac{1}{\Im m_\pm(z, \omega) + \frac{1}{2} \Im z} \right) \\
&= \frac{1}{\Im z} \mathbb{E} \left(\frac{\frac{\Im z}{\Im m_\pm(z, \omega)}}{1 + \frac{\Im z}{2 \Im m_\pm(z, \omega)}} \right) \\
&\leq \frac{1}{\Im z} \mathbb{E} \left(\log \left(1 + \frac{\Im z}{\Im m_\pm(z, \omega)} \right) \right) \\
&= \frac{2\gamma(z)}{\Im z},
\end{aligned}$$

which is (17). We used (19) in the third step and Proposition 2.1 in the last step.

Notice that $n_+(z, \omega) + n_-(z, \omega) = \Im b(z, \omega)$. Thus the integrand on the left-hand side of (18) is equal to

$$\begin{aligned} \frac{\left[\frac{1}{n_+} + \frac{1}{n_-} \right] \cdot [(n_+ + n_-)^2 - 4n_+n_- + (\Re b)^2]}{|b|^2} &= \frac{\left[\frac{1}{n_+} + \frac{1}{n_-} \right] \cdot [|b|^2 - 4n_+n_-]}{|b|^2} \\ &= \frac{1}{n_+} + \frac{1}{n_-} - 4 \frac{n_+ + n_-}{|b|^2} \\ &= \frac{1}{n_+} + \frac{1}{n_-} + 4 \Im \left(\frac{1}{b} \right). \end{aligned}$$

The bound (18) now follows from (17) and Proposition 2.2. \square

Proof of the Kotani half of Theorem 4. The Thouless formula (4) says that

$$\gamma(z) = \int \log |z - E'| dk(E') = \Re \int \log(z - E') dk(E')$$

and hence $-\gamma(z)$ is the real part of a function whose derivative is a Borel transform (namely, of the measure dk). By general properties of the Borel transform, it follows that the limit $\gamma'(E + i0)$ exists for Lebesgue almost every $E \in \mathbb{R}$ and, in particular, for almost every $E \in \mathcal{Z}$. For these E , we have that

$$(20) \quad \lim_{\varepsilon \downarrow 0} \frac{\gamma(E + i\varepsilon)}{\varepsilon} = \lim_{\varepsilon \downarrow 0} \frac{\gamma(E + i\varepsilon) - \gamma(E)}{\varepsilon - 0} = \lim_{\varepsilon \downarrow 0} \frac{\partial \gamma}{\partial (\Im z)}(E + i\varepsilon),$$

and in particular, the limit is finite. Thus, by (17),

$$\limsup_{\varepsilon \downarrow 0} \mathbb{E} \left(\frac{1}{\Im m_{\pm}(E + i\varepsilon, \omega)} \right) < \infty$$

for almost every $E \in \mathcal{Z}$. Since m_{\pm} are Borel transforms as well (of the spectral measures associated with half-line restrictions of H_{ω}), we also have that, for every $\omega \in \Omega$, $m_{\pm}(E + i0, \omega)$ exists for Lebesgue almost every $E \in \mathbb{R}$, and hence, for almost every E , $m_{\pm}(E + i0, \omega)$ exists for almost every ω . Combining the last two observations with Fatou's lemma, we find that

$$(21) \quad \mathbb{E} \left(\frac{1}{\Im m_{\pm}(E + i0, \omega)} \right) < \infty$$

for almost every E in \mathcal{Z} . So, for almost every $\omega \in \Omega$ and $E \in \mathcal{Z}$, $\Im m_{\pm}(E + i0, \omega) > 0$.

On the other hand, $m_+(E + i\varepsilon, \omega) + m_-(E + i\varepsilon, \omega) + E + i\varepsilon - V_{\omega}(0)$ has a finite limit for almost every $\omega \in \Omega$ and $E \in \mathcal{Z}$.

Hence, (15) shows that $0 < \Im G_{\omega}(0, 0; E + i0) < \infty$ for almost every $\omega \in \Omega$ and $E \in \mathcal{Z}$, which implies the result. \square

Denote the measure associated with the Herglotz function $G_{\omega}(0, 0; z)$ by ν_{ω} , that is,

$$G_{\omega}(0, 0; z) = \int \frac{d\nu_{\omega}(E)}{E - z}.$$

The results above imply the following for μ -almost every ω :

$$\begin{aligned} \nu_{\omega}^{(\text{ac})}(E) &= 0 \text{ for Lebesgue almost every } E \in \mathbb{R} \setminus \mathcal{Z}, \\ \nu_{\omega}^{(\text{ac})}(E) &> 0 \text{ for Lebesgue almost every } E \in \mathcal{Z}. \end{aligned}$$

Here, $\nu_\omega^{(\text{ac})}(E)$ denotes the density of the absolutely continuous part of ν_ω . Write $k^{(\text{ac})}(E)$ for the density of the absolutely continuous part of the density of states measure.

There is a direct relation between these densities [47]:

Theorem 5 (Kotani 1997). *For almost every $E \in \mathcal{Z}$,*

$$(22) \quad k^{(\text{ac})}(E) = \mathbb{E} \left(\nu_\omega^{(\text{ac})}(E) \right).$$

Proof. The inequality “ \geq ” in (22) follows from (2) (i.e., the density of states measure is the average of the measures ν_ω) and the fact that the average of absolutely continuous measures is absolutely continuous.

To prove the opposite inequality, we first note that for almost every $E \in \mathcal{Z}$, (16), (20), and Cauchy-Riemann imply

$$(23) \quad k^{(\text{ac})}(E) = \frac{1}{\pi} \lim_{\varepsilon \downarrow 0} \frac{\gamma(E + i\varepsilon)}{\varepsilon}.$$

Because of (20), (21) and Fatou’s lemma, (18) implies that for almost every pair $(E, \omega) \in \mathcal{Z} \times \Omega$,

$$(24) \quad \Im m_+(E + i0, \omega) = \Im m_-(E + i0, \omega)$$

and

$$(25) \quad \Re m_+(E + i0, \omega) + \Re m_-(E + i0, \omega) + E - V_\omega(0) = 0.$$

Thus, for almost every $(E, \omega) \in \mathcal{Z} \times \Omega$,

$$(26) \quad \begin{aligned} \nu_\omega^{(\text{ac})}(E) &= \frac{1}{\pi} \Im G_\omega(0, 0; E + i0) \\ &= \frac{1}{\pi} \Im \frac{-1}{m_+(E + i0, \omega) + m_-(E + i0, \omega) + E - V_\omega(0)} \\ &= \frac{1}{\pi} \Im \frac{-1}{2i \Im m_+(E + i0, \omega)} \\ &= \frac{1}{2\pi} \frac{1}{\Im m_+(E + i0, \omega)} \end{aligned}$$

Let P_ε be the Poisson kernel for the upper half-plane, that is,

$$P_\varepsilon(E) = \frac{1}{\pi} \frac{\varepsilon}{E^2 + \varepsilon^2}.$$

Write

$$C_\varepsilon(E) = \int_{\mathcal{Z}} P_\varepsilon(E - E') dE'$$

and

$$\tilde{P}_\varepsilon(E, E') = P_\varepsilon(E - E') C_\varepsilon(E)^{-1}.$$

Then, by (26) and Jensen's inequality, we obtain for almost every $(E, \omega) \in \mathcal{Z} \times \Omega$,

$$\begin{aligned}
\int_{\mathbb{R}} \nu_{\omega}^{(\text{ac})}(E') P_{\varepsilon}(E - E') dE' &\geq \int_{\mathcal{Z}} \nu_{\omega}^{(\text{ac})}(E') P_{\varepsilon}(E - E') dE' \\
&= \int_{\mathcal{Z}} \left(\frac{1}{2\pi} \frac{1}{\Im m_{+}(E' + i0, \omega)} \right) P_{\varepsilon}(E - E') dE' \\
&= C_{\varepsilon}(E) \int_{\mathcal{Z}} \left(\frac{1}{2\pi} \frac{1}{\Im m_{+}(E' + i0, \omega)} \right) \tilde{P}_{\varepsilon}(E, E') dE' \\
&\geq C_{\varepsilon}(E) \left(\int_{\mathcal{Z}} \left(\frac{1}{2\pi} \frac{1}{\Im m_{+}(E' + i0, \omega)} \right)^{-1} \tilde{P}_{\varepsilon}(E, E') dE' \right)^{-1} \\
&\geq C_{\varepsilon}(E)^2 \left(\int_{\mathbb{R}} \left(\frac{1}{2\pi} \frac{1}{\Im m_{+}(E' + i0, \omega)} \right)^{-1} P_{\varepsilon}(E - E') dE' \right)^{-1} \\
&= \frac{C_{\varepsilon}(E)^2}{2\pi} \frac{1}{\Im m_{+}(E + i\varepsilon, \omega)}
\end{aligned}$$

Thus, for almost every $E \in \mathcal{Z}$,

$$\int_{\mathbb{R}} \mathbb{E} \left(\nu_{\omega}^{(\text{ac})}(E') \right) P_{\varepsilon}(E - E') dE' \geq \frac{C_{\varepsilon}(E)^2}{2\pi} \mathbb{E} \left(\frac{1}{\Im m_{+}(E + i\varepsilon, \omega)} \right),$$

and hence

$$(27) \quad \mathbb{E} \left(\nu_{\omega}^{(\text{ac})}(E) \right) \geq \frac{1}{2\pi} \limsup_{\varepsilon \downarrow 0} \mathbb{E} \left(\frac{1}{\Im m_{+}(E + i\varepsilon, \omega)} \right)$$

since $C_{\varepsilon}(E) < 1$ and $C_{\varepsilon}(E) \rightarrow 1$ as $\varepsilon \downarrow 0$.

Using (23), Proposition 2.1, the inequality $\log(1+x) \leq x$ for $x \geq 0$, and then (27), we find that

$$\begin{aligned}
k^{(\text{ac})}(E) &= \frac{1}{\pi} \lim_{\varepsilon \downarrow 0} \frac{\gamma(E + i\varepsilon)}{\varepsilon} \\
&= \frac{1}{\pi} \lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \mathbb{E} \left(\log \left(1 + \frac{\varepsilon}{\Im m_{\pm}(E + i\varepsilon, \omega)} \right) \right) \\
&\leq \frac{1}{2\pi} \limsup_{\varepsilon \downarrow 0} \mathbb{E} \left(\frac{1}{\Im m_{\pm}(E + i\varepsilon, \omega)} \right) \\
&\leq \mathbb{E} \left(\nu_{\omega}^{(\text{ac})}(E) \right),
\end{aligned}$$

concluding the proof of “ \leq ” in (22). \square

Corollary 1. *The spectrum is almost surely purely absolutely continuous if and only if the integrated density of states is absolutely continuous and the Lyapunov exponent vanishes almost everywhere with respect to the density of states measure.*

There is a different approach to purely absolutely continuous spectrum as pointed out by Yoram Last (unpublished):⁴ Using a result of Deift and Simon [30, Theorem 7.1], one can show that there is a set $\mathcal{R} \subseteq \mathcal{Z}$ such that $\text{Leb}(\mathcal{Z} \setminus \mathcal{R}) = 0$ and \mathcal{R} has zero singular spectral measure for every $\omega \in \Omega$ due to the absence of subordinate solutions.

⁴The author is grateful to Barry Simon for bringing this to his attention.

The spectrum naturally breaks up into the two components \mathcal{Z} and $\Sigma \setminus \mathcal{Z}$. It is known that either set can support singular continuous spectrum as demonstrated by the almost Mathieu operator at critical and super-critical coupling. However, as we have seen, $\Sigma \setminus \mathcal{Z}$ does not support any absolutely continuous part of the spectral measures. Trivially, Anderson localization is impossible in \mathcal{Z} . However, it is tempting to expect even more:

Problem 1. *Prove or disprove that for all ergodic families, the operators H_ω have no eigenvalues in \mathcal{Z} .*

If the answer is affirmative, this will in particular imply the absence of eigenvalues in a number of special cases, such as the operators considered in Section 6 and the critical almost Mathieu operator.⁵

3. THE INDUCED MEASURE AND ITS TOPOLOGICAL SUPPORT

Fix a compact subset R of \mathbb{R} . Endow $R^{\mathbb{Z}}$ with product topology, which makes it a compact metric space. If $V \in R^{\mathbb{Z}}$, we define the functions m_\pm by

$$m_\pm(z) = \mp \frac{u_\pm(1)}{u_\pm(0)},$$

where u_\pm solves

$$(28) \quad u(n+1) + u(n-1) + V(n)u(n) = zu(n)$$

and is ℓ^2 at $\pm\infty$.

Denote $\mathbb{Z}_+ = \{1, 2, 3, \dots\}$ and $\mathbb{Z}_- = \{0, -1, -2, \dots\}$. It is well known that the maps $\mathcal{M}_\pm : V_\pm = V|_{\mathbb{Z}_\pm} \mapsto m_\pm$ are one-one and continuous with respect to uniform convergence on compacta on the m -function side. The m -functions m_\pm are Herglotz functions, they have boundary values almost everywhere on the real axis, and they are completely determined by their boundary values on any set of positive Lebesgue measure.

We will be interested in those V for which the functions m_+, m_- obey identities like (24) and (25), that is,

$$(29) \quad m_-(E + i0) = -\overline{m_+(E + i0)}$$

for a rich set of energies. Thus, for a set $\mathcal{Z} \subseteq \mathbb{R}$, we let

$$\mathcal{D}(\mathcal{Z}) = \{V \in R^{\mathbb{Z}} : m_\pm \text{ associated with } V \text{ obey (29) for a.e. } E \in \mathcal{Z}\}.$$

On $R^{\mathbb{Z}}$, define the shift transformation $[S(V)](n) = V(n+1)$.

Lemma 3.1. *Suppose that $\mathcal{Z} \subset \mathbb{R}$ has positive Lebesgue measure. Then:*

- (a) $\mathcal{D}(\mathcal{Z})$ is S -invariant and closed in $R^{\mathbb{Z}}$.
- (b) For $V \in \mathcal{D}(\mathcal{Z})$, denote the restrictions to \mathbb{Z}_\pm by V_\pm . Then V_- determines V_+ uniquely among elements of $\mathcal{D}(\mathcal{Z})$ and vice versa.
- (c) If there exist $V^{(m)}, V \in \mathcal{D}(\mathcal{Z})$ such that $V_-^{(m)} \rightarrow V_-$ pointwise, then $V_+^{(m)} \rightarrow V_+$ pointwise.

⁵For this particular model, this would provide an alternative to the proof based on self-duality and zero-measure spectrum.

Proof. (a) If u_1, u_2 denote the solutions of (28) that obey $u_1(0) = u_2(1) = 1$ and $u_1(1) = u_2(0) = 0$, then we can write (note that we may normalize u_{\pm} by $u_{\pm}(0) = 1$)

$$u_{\pm}(n) = u_1(n) \mp m_{\pm}(z)u_2(n).$$

Let us denote the m -functions associated with $S(V)$ by \tilde{m}_{\pm} . Clearly,

$$\tilde{m}_{\pm}(z) = \mp \frac{u_1(2) \mp m_{\pm}(z)u_2(2)}{u_1(1) \mp m_{\pm}(z)u_2(1)}.$$

Since the $u_j(m)$ are polynomials in z with real coefficients, this shows that

$$\tilde{m}_{-}(E + i0) = -\overline{\tilde{m}_{+}(E + i0)}$$

for almost every $E \in \mathcal{Z}$ and hence $\mathcal{D}(\mathcal{Z})$ is S -invariant. It follows from the continuity of the maps \mathcal{M}_{\pm} that $\mathcal{D}(\mathcal{Z})$ is closed. (For a proof that the identity between the boundary values of the associated m -functions is preserved after taking limits, see [44, Lemma 5] and [45, Lemma 7.4].)

(b) V_{-} determines m_{-} and then (29) determines the boundary values of m_{+} on a set of positive Lebesgue measure. By general properties of Herglotz functions, this determines m_{+} (and hence V_{+}) completely. By the same argument, V_{+} determines V_{-} .

(c) By compactness, there is a subsequence of $\{V^{(m)}\}$ that converges pointwise, that is, there is \tilde{V} such that $V^{(m_k)} \rightarrow \tilde{V}$ as $k \rightarrow \infty$. By part (a), $\tilde{V} \in \mathcal{D}(\mathcal{Z})$. By assumption, $V_{-} = \tilde{V}_{-}$. Thus, by part (b), $V_{+} = \tilde{V}_{+}$, and hence $V = \tilde{V}$. Consequently, $V_{+}^{(m_k)} \rightarrow V_{+}$ pointwise. In fact, we claim that $V_{+}^{(m)} \rightarrow V_{+}$ pointwise. Otherwise, we could reverse the argument (i.e., go from right to left) and show that $V_{-}^{(\tilde{m}_k)} \not\rightarrow V_{-}$ for some other subsequence. \square

We will now derive two important consequences of Lemma 3.1: the absence of absolutely continuous spectrum for topologically non-deterministic families and the support theorem. To do so, we will consider the push-forward ν of μ on the sequence space and its topological support.

More precisely, given an ergodic dynamical system (Ω, μ, T) and a measurable bounded sampling function $f : \Omega \rightarrow \mathbb{R}$ defining potentials $V_{\omega}(n) = f(T^n \omega)$ as before, we associate the following dynamical system $(R^{\mathbb{Z}}, \nu, S)$: R is a compact set that contains the range of f , ν is the Borel measure on $R^{\mathbb{Z}}$ induced by μ via $\Phi(\omega) = V_{\omega}$ (i.e., $\nu(A) = \mu(\Phi^{-1}(A))$), and S is the shift transformation on $R^{\mathbb{Z}}$ introduced above. Recall that the *topological support* of ν , $\text{supp } \nu$, is given by the intersection of all compact sets $B \subseteq R^{\mathbb{Z}}$ with $\nu(B) = 1$. Clearly, $\text{supp } \nu$ is closed and S -invariant.

Theorem 6 (Kotani 1989⁶). *Let (Ω, μ, T, f) and $(R^{\mathbb{Z}}, d\nu, S)$ be as just described. Assume that the set*

$$\mathcal{Z} = \{E \in \mathbb{R} : \gamma(E) = 0\}.$$

has positive Lebesgue measure. Then,

- (a) *Each $V \in \text{supp } \nu$ is determined completely by V_{-} (resp., V_{+}).*
- (b) *If we let*

$$(\text{supp } \nu)_{\pm} = \{V_{\pm} : V \in \text{supp } \nu\},$$

⁶The result appears explicitly in the 1989 paper [46]. The main ingredients of the proof, however, were found earlier [44, 45].

then the mappings

$$(30) \quad E_{\pm} : (\text{supp } \nu)_{\pm} \ni V_{\pm} \mapsto V_{\mp} \in (\text{supp } \nu)_{\mp}$$

are continuous with respect to pointwise convergence.

Proof. (a) By our earlier results, we know that $\mathcal{D}(\mathcal{Z})$ is compact and has full ν -measure. Thus, $\text{supp } \nu \subseteq \mathcal{D}(\mathcal{Z})$ and the assertion follows from Lemma 3.1.(b).

(b) This follows from Lemma 3.1.(c). \square

We say that (Ω, μ, T, f) is *topologically deterministic* if there exist continuous mappings $E_{\pm} : (\text{supp } \nu)_{\pm} \rightarrow (\text{supp } \nu)_{\mp}$ that are formal inverses of one another and obey $V_{-}^{\#} \in \text{supp } \nu$ for every $V_{-} \in (\text{supp } \nu)_{-}$, where

$$V_{-}^{\#}(n) = \begin{cases} V_{-}(n) & n \leq 0, \\ E_{-}(V_{-})(n) & n \geq 1. \end{cases}$$

This also implies $V_{+}^{\#} \in \text{supp } \nu$ for every $V_{+} \in (\text{supp } \nu)_{+}$, where

$$V_{+}^{\#}(n) = \begin{cases} V_{+}(n) & n \geq 1, \\ E_{+}(V_{+})(n) & n \leq 0. \end{cases}$$

Otherwise, (Ω, μ, T, f) is *topologically non-deterministic*.

Corollary 2. *If (Ω, μ, T, f) is topologically non-deterministic, then $\Sigma_{\text{ac}} = \emptyset$.*

Our next application of Lemma 3.1 is the so-called support theorem; compare [44]. For a Borel measure ν on $R^{\mathbb{Z}}$, let $\Sigma_{\text{ac}}(\nu) \subseteq \mathbb{R}$ denote the almost sure absolutely continuous spectrum, that is, $\sigma_{\text{ac}}(\Delta + V) = \Sigma_{\text{ac}}(\nu)$ for ν almost every V . If ν comes from (Ω, μ, T, f) , then $\Sigma_{\text{ac}}(\nu)$ coincides with the set Σ_{ac} introduced earlier. The support theorem says that $\Sigma_{\text{ac}}(\nu)$ is monotonically decreasing in the support of ν .

Theorem 7 (Kotani 1985). *For every $V \in \text{supp } \nu$, we have $\sigma_{\text{ac}}(\Delta + V) \supseteq \Sigma_{\text{ac}}(\nu)$. In particular, $\text{supp } \nu_1 \subseteq \text{supp } \nu_2$ implies that $\Sigma_{\text{ac}}(\nu_1) \supseteq \Sigma_{\text{ac}}(\nu_2)$.*

Proof. We know that

$$\text{supp } \nu \subseteq \mathcal{D}(\mathcal{Z}) = \{V \in R^{\mathbb{Z}} : m_{\pm} \text{ associated with } V \text{ obey (29) for a.e. } E \in \mathcal{Z}\}.$$

Bearing in mind the Riccati equation (13), a calculation like the one in (26) therefore shows that for every $V \in \text{supp } \nu$, the Green function associated with the operator $\Delta + V$ obeys $\Im G(0, 0; E + i0) > 0$ for almost every $E \in \mathcal{Z}$. This implies $\overline{\mathcal{Z}}^{\text{ess}} \subseteq \sigma_{\text{ac}}(\Delta + V)$ and hence the result by Theorem 4. \square

A different proof may be found in Last-Simon [51, Sect. 6]. Here is a typical application of the support theorem:

Corollary 3. *Let Per_{ν} be the set of $V \in \text{supp } \nu$ that are periodic, that is, $S^p V = V$ for some $p \in \mathbb{Z}_{+}$. Then,*

$$\Sigma_{\text{ac}}(\nu) \subseteq \bigcap_{V \in \text{Per}_{\nu}} \sigma(\Delta + V).$$

If there are sufficiently many gaps in the spectra of these periodic operators, one can show in this way that $\Sigma_{\text{ac}}(\nu)$ is empty.

Corollaries 2 and 3 have been used in a variety of scenarios to prove the absence of absolutely continuous spectrum. In fact, while the Kotani half of Theorem 4 concerns the *presence* of absolutely continuous spectrum on \mathcal{Z} , it could be argued

that the criteria for the *absence* of absolutely continuous spectrum that are by-products of the theory have been applied more often. This is to a certain extent due to the fact that the majority of the ergodic families of Schrödinger operators are expected to have no absolutely continuous spectrum. The following “conjecture” is tempting because it is supported by a plethora of results, both on the positive side and on the negative side. It has been verbally suggested by Yoram Last and it has appeared explicitly in print in several places, including [39, 48].

Problem 2. *Show that $\text{Leb}(\mathcal{Z}) > 0$ implies almost periodicity, that is, the closure in $\ell^\infty(\mathbb{Z})$ of the set of translates of V_ω is compact.*

Namely, the presence of (purely) absolutely continuous spectrum is known for all periodic potentials, many limit-periodic potentials,⁷ and some quasi-periodic potentials (that are all uniformly almost periodic). On the other hand, the absence of absolutely continuous spectrum is known for large classes of non-almost periodic ergodic potentials. We will see some instances of the latter statement below. Nevertheless, proving this conjecture is presumably very hard and it would already be interesting to find further specific results that support the conjecture. For example, it is an open (and seemingly hard) problem to prove the absence of absolutely continuous spectrum for potentials defined by the skew shift.

We close this section with a problem concerning strips. As was mentioned at the end of the Introduction, Kotani and Simon developed the analog of Kotani theory for discrete and continuous Schrödinger operators with matrix-valued potentials in their 1988 paper [49]. This framework includes in particular discrete Schrödinger operators on strips. That is, operators of the form $\Delta + V_\omega$ on $\ell^2(\mathbb{Z} \times \{1, \dots, L\})$, where Δ is again given by the summation over nearest neighbors. Transfer matrices are now $2L \times 2L$ and, modulo symmetry, there are L Lyapunov exponents, $\gamma_L(E) \geq \gamma_{L-1}(E) \geq \dots \geq \gamma_1(E) \geq 0$. For the general matrix-valued situation, they proved that the largest Lyapunov exponent is positive for almost every energy if the potential is non-deterministic. This result cannot be improved in this general setting. However, it is reasonable to expect that for strips, the following result should hold.

Problem 3. *Prove that for non-deterministic Schrödinger operators on a strip, all Lyapunov exponents are non-zero for Lebesgue almost all energies.*

4. POTENTIALS GENERATED BY THE DOUBLING MAP

In this section we discuss potentials defined over the doubling map, that is, Example (c) from the introduction. The underlying dynamical system is strongly mixing and one would hope that the spectral theory of the associated operators is akin to that of the Anderson model, where the potentials are generated by independent, identically distributed random variables. Alas, by dropping independence one loses the availability of most tools that have proven useful in the study of the Anderson model.

While localization is expected for Schrödinger operators with potentials over the doubling map, it has not been shown to hold in reasonable generality. There are only two localization results in the literature, and each of them is to some

⁷A sequence V is limit-periodic if there are periodic sequences $V^{(m)}$ such that $\|V - V^{(m)}\|_\infty \rightarrow 0$ as $m \rightarrow \infty$. See, for example, Avron-Simon [7], Chulaevsky [20], Chulaevsky-Molchanov [21], and Pastur-Tkachenko [58].

extent unsatisfactory. The first result was found by Bourgain and Schlag [17], who proved localization at small coupling and away from small intervals about the energies ± 2 and 0. Both assumptions seem unnatural. The other result is due to Damanik and Killip [27], who proved localization for essentially all f but only for Lebesgue almost every boundary condition at the origin (recall that we are dealing with operators in $\ell^2(\mathbb{Z}_+)$). A result holding for fixed boundary condition would of course be more desirable. To this end, Damanik and Killip were at least able to show the absence of absolutely continuous spectrum for fixed boundary condition in complete generality. These results are indeed immediate consequences of Kotani theory and spectral averaging and we give the short proofs below for the reader's convenience.

The first step in a localization proof for a one-dimensional Schrödinger operator is typically a proof of positive Lyapunov exponents for many energies. For the Anderson model, this can be done for all energies using Fürstenberg's theorem or for Lebesgue almost all energies using Kotani theory. At small coupling there is also a perturbative approach due to Pastur and Figotin [57]. The extension of the approach based on Fürstenberg's theorem to potentials generated by the doubling map is not obvious; see, however, [4]. The perturbative approach extends quite nicely as shown by Chulaevsky and Spencer [22]. Their results form the basis for the proof of the partial localization result in [17]. Finally, the approach based on Kotani theory also extends as we will now explain.

Theorem 8 (Damanik-Killip 2005). *Suppose that $f \in L^\infty(\mathbb{T})$ is non-constant and $V_\omega(n) = f(2^n \omega)$ for $n \geq 1$. Then, the Lyapunov exponent $\gamma(E)$ is positive for Lebesgue almost every $E \in \mathbb{R}$ and the absolutely continuous spectrum of the operator H_ω in $\ell^2(\mathbb{Z}_+)$ is empty for Lebesgue almost every $\omega \in \mathbb{T}$.*

Proof. Since the proof of this result is so short, we reproduce it here in its entirety. The first step is to conjugate the doubling map T to a symbolic shift via the binary expansion. Let $\tilde{\Omega}_+ = \{0, 1\}^{\mathbb{Z}_+}$ and define $D : \tilde{\Omega}_+ \rightarrow \mathbb{T}$ by $D(\omega) = \sum_{n=1}^{\infty} \omega_n 2^{-n}$. The shift transformation, $S : \tilde{\Omega}_+ \rightarrow \tilde{\Omega}_+$, is given by $(S\tilde{\omega})_n = \tilde{\omega}_{n+1}$. Clearly, $D \circ S = T \circ D$.

Next we introduce a family of whole-line operators as follows. Let $\tilde{\Omega} = \{0, 1\}^{\mathbb{Z}}$ and define, for $\tilde{\omega} \in \tilde{\Omega}$, the operator

$$[H_{\tilde{\omega}}\phi](n) = \phi(n+1) + \phi(n-1) + V_{\tilde{\omega}}(n)\phi(n)$$

in $\ell^2(\mathbb{Z})$, where

$$V_{\tilde{\omega}}(n) = f[D(\{\tilde{\omega}_n, \tilde{\omega}_{n+1}, \tilde{\omega}_{n+2}, \dots\})].$$

The family $\{H_{\tilde{\omega}}\}_{\tilde{\omega} \in \tilde{\Omega}}$ is non-deterministic since $V_{\tilde{\omega}}$ restricted to \mathbb{Z}_+ only depends on $\{\tilde{\omega}_n\}_{n \geq 1}$ and hence, by non-constancy of f , we cannot determine the values of $V_{\tilde{\omega}}(n)$ for $n \leq 0$ uniquely from the knowledge of $V_{\tilde{\omega}}(n)$ for $n \geq 1$. It follows from Corollary 2 that the Lyapunov exponent for $\{H_{\tilde{\omega}}\}_{\tilde{\omega} \in \tilde{\Omega}}$ is almost everywhere positive and $\sigma_{ac}(H_{\tilde{\omega}})$ is empty for almost every $\tilde{\omega} \in \tilde{\Omega}$ with respect to the $(\frac{1}{2}, \frac{1}{2})$ -Bernoulli measure on $\tilde{\Omega}$.

Finally, let us consider the restrictions of $H_{\tilde{\omega}}$ to $\ell^2(\mathbb{Z}_+)$, that is, let $H_{\tilde{\omega}}^+ = E^* H_{\tilde{\omega}} E$, where $E : \ell^2(\mathbb{Z}_+) \rightarrow \ell^2(\mathbb{Z})$ is the natural embedding. Observe that $H_{\tilde{\omega}}^+ = H_\omega$, where $\omega = D(\{\tilde{\omega}_1, \tilde{\omega}_2, \tilde{\omega}_3, \dots\})$. This immediately implies the statement on the positivity of the Lyapunov exponent for the family $\{H_\omega\}_{\omega \in \mathbb{T}}$. As finite-rank

perturbations preserve absolutely continuous spectrum, $\sigma_{\text{ac}}(H_{\tilde{\omega}}^+) \subseteq \sigma_{\text{ac}}(H_{\tilde{\omega}})$ for every $\tilde{\omega} \in \tilde{\Omega}$. This proves that $\sigma_{\text{ac}}(H_{\tilde{\omega}}^+) = \emptyset$ for almost every $\tilde{\omega} \in \tilde{\Omega}$. \square

Given $\phi \in (-\pi/2, \pi/2)$, let $H_{\omega}^{(\phi)}$ denote the operator which acts on $\ell^2(\mathbb{Z}_+)$ as in (1), but with $\psi(0)$ given by $\cos(\phi)\psi(0) + \sin(\phi)\psi(1) = 0$. Thus, the original operator (with a Dirichlet boundary condition) corresponds to $\phi = 0$. Theorem 8 implies the following result for this family of operators:

Corollary 4. *Suppose that f is measurable, bounded, and non-constant. Then, for almost every $\phi \in (-\pi/2, \pi/2)$ and almost every $\omega \in \mathbb{T}$, the operator $H_{\omega}^{(\phi)}$ in $\ell^2(\mathbb{Z}_+)$ with potential $V_{\omega}(n) = f(2^n \omega)$ has pure point spectrum and all eigenfunctions decay exponentially at infinity.*

Proof. This is standard and follows quickly from spectral averaging; see, for example, [57, Theorem 13.4] or [61, Section 12.3]. \square

For a localization proof without the need for spectral averaging, it will be necessary to prove the positivity of the Lyapunov exponent for a larger set of energies. Sufficient, for example, is positivity away from a discrete set of exceptional energies. For moderately small coupling, such a result will be contained in [4]. The problem for other values of the coupling constant is still open.

Problem 4. *Find a class of functions $f \in L^{\infty}(\mathbb{T})$ such that for every $\lambda \neq 0$, the Lyapunov exponent associated with the potentials $V_{\omega}(n) = \lambda f(2^n \omega)$ is positive away from a (λ -dependent) discrete set of energies.*

In connection with this problem, important obstructions have been found by Bochi [10] and Bochi and Viana [11]. Namely, positivity of $\gamma(E)$ away from a discrete set will fail generically in $C(\mathbb{T})$ (this result holds for rather general underlying dynamics) and hence the Hölder continuity assumptions made in [4] and [22] are natural.

In some sense a large value of λ alone should ensure the positivity of the Lyapunov exponent. This is indeed the basis of several results for quasi-periodic potentials or potentials generated by the skew-shift. For hyperbolic base transformations such as the doubling map, however, there is a competition between two different kinds of hyperbolic behavior that presents problems that have not been solved yet. Moreover, in the large coupling regime, it would be especially interesting to prove uniform (in energy) lower bounds on the Lyapunov exponents along with the natural $\log \lambda$ asymptotics.

Problem 5. *Find a class of functions $f \in L^{\infty}(\mathbb{T})$ such that for every $\lambda \geq \lambda_0(g)$, the Lyapunov exponent associated with the potentials $V_{\omega}(n) = \lambda f(2^n \omega)$ obeys $\inf_E \gamma(E) \geq c \log \lambda$ for some suitable positive constant c .*

In this context it should be noted that Herman's subharmonicity proof for trigonometric polynomials over ergodic shifts on the torus, [36], works in the case of the doubling map.⁸ It seems much less clear, however, how to carry over the Sorets-Spencer proof for real-analytic f , [62], from the case of irrational rotations of the circle to the case of the doubling map, let alone the proof of Bourgain for real-analytic functions over ergodic shifts on higher-dimensional tori [14].

⁸The author is grateful to Kristian Bjerklöv for pointing this out.

For $\mathrm{SL}(2, \mathbb{R})$ cocycles over the doubling map that are not of Schrödinger form (i.e., with a general $\mathrm{SL}(2, \mathbb{R})$ matrix replacing (3)), Young has developed a method for proving positive Lyapunov exponents “at large coupling” that works in the C^1 category [64]. Her method does not immediately apply to Schrödinger cocycles, but it would be interesting to find a suitable extension.

Problem 6. *Modify Young’s method and apply it to Schrödinger cocycles.*

5. ABSENCE OF AC SPECTRUM FOR ROUGH POTENTIALS

It is in some way surprising that a rough sampling function f can make the resulting potentials non-deterministic. Traditionally, non-determinism had been thought of as a feature induced by the underlying dynamics. In particular, quasi-periodic potentials (in the generalized sense considered in this paper, which allows discontinuous f ’s) had for a long time been considered deterministic. The situation changed with an important observation by Kotani in his short 1989 paper [46].

He proved the following very general result:

Theorem 9 (Kotani 1989). *Suppose that (Ω, T, μ) is ergodic, $f : \Omega \rightarrow \mathbb{R}$ takes finitely many values, and the resulting potentials V_ω are μ -almost surely not periodic. Then, $\mathrm{Leb}(\mathcal{Z}) = 0$ and therefore $\Sigma_{\mathrm{ac}} = \emptyset$.*

In particular, operators with quasi-periodic potentials of the form

$$V_\omega(n) = \lambda \sum_{m=1}^N \gamma_m \chi_{[a_{m-1}, a_m)}(n\alpha + \omega),$$

where $0 = a_0 < a_1 < \cdots < a_{N-1} < a_N = 1$, $\gamma_1, \dots, \gamma_N \in \mathbb{R}$ (taking at least two values) and $\lambda \neq 0$, have no absolutely continuous spectrum. Note that this result holds for all non-zero couplings and hence it is particularly surprising for small values of λ . We will have more to say about these potentials in the next section.

The family $\{V_\omega\}_{\omega \in \mathbb{T}}$ does not seem to be non-deterministic in an intuitive sense as ω is uniquely determined by the sequence $V_\omega|_{\mathbb{Z}_-}$. However, the family becomes non-deterministic when we pass to the closed topological support of the induced measure on $\{\lambda\gamma_1, \dots, \lambda\gamma_N\}^{\mathbb{Z}}$. Then we will indeed find two distinct sequences that belong to the support of the induced measure, whose restrictions to \mathbb{Z}_- coincide. This fact will become more transparent when we discuss Theorem 10 below. Kotani’s proof proceeded in a slightly different way; he proved that there can be no *continuous* mapping from the left half-line to the right half-line.

Proof of Theorem 9. As above we denote the push-forward of μ under the map $\omega \mapsto V_\omega$ by ν . It suffices to show that $\mathrm{Leb}(\mathcal{Z}) > 0$ implies that $\mathrm{supp} \nu$ is finite since then all elements of $\mathrm{supp} \nu$ are periodic.

By the continuity of the maps E_\pm from (30) and the fact that $\mathrm{Ran} f$ is finite, there is a finite M such that the knowledge of $V(-M), \dots, V(-1)$ determines $V(0)$ uniquely for $V \in \mathrm{supp} \nu$. Now shift and iterate! It follows that $V(-M), \dots, V(-1)$ completely determine $\{V(n) : n \geq 0\}$ and hence $\mathrm{supp} \nu$ has cardinality at most $(\#\mathrm{Ran} f)^M$. \square

Consider the case where Ω is a compact metric space, T is a homeomorphism, and μ is an ergodic Borel probability measure. This covers most, if not all, applications of interest. In this scenario, Damanik and Killip realized in [26] that finite range of f is not essential. What is important, however, is that f is discontinuous at some

point $\omega_0 \in \Omega$. One can then use this point of discontinuity to actually “construct” two elements of $\text{supp } \nu$ that coincide on a half-line.

We say that $l \in \mathbb{R}$ is an *essential limit* of f at ω_0 if there exists a sequence $\{\Omega_k\}$ of sets each of positive measure such that for any sequence $\{\omega_k\}$ with $\omega_k \in \Omega_k$, both $\omega_k \rightarrow \omega_0$ and $f(\omega_k) \rightarrow l$. If f has more than one essential limit at ω_0 , we say that f is *essentially discontinuous* at this point.

Theorem 10 (Damanik-Killip 2005). *Suppose Ω is a compact metric space, $T : \Omega \rightarrow \Omega$ a homeomorphism, and μ an ergodic Borel probability measure. If there is an $\omega_0 \in \Omega$ such that f is essentially discontinuous at ω_0 but continuous at all points $T^n \omega_0$, $n < 0$, then $\Sigma_{ac} = \emptyset$.*

Proof. We again denote the induced measure by ν . For each essential limit l of f at ω_0 , we will find $V_l \in \text{supp } \nu$ with $V_l(0) = l$. By assumption and construction, $V_l(n)$ is independent of l for every $n < 0$. This shows $\text{Leb}(\mathcal{Z}) = 0$ and hence $\Sigma_{ac} = \emptyset$.

Let l be an essential limit of f at ω_0 and let $\{\Omega_k\}$ be a sequence of sets which exhibits the fact that l is an essential limit of f . Since each has positive μ -measure, we can find points $\omega_k \in \Omega_k$ so that V_{ω_k} is in $\text{supp } \nu$; indeed, this is the case for almost every point in Ω_k .

As $\omega_k \rightarrow \omega_0$ and f is continuous at each of the points $T^n \omega_0$, $n < 0$, it follows that $V_{\omega_k}(n) \rightarrow V_{\omega_0}(n)$ for each $n < 0$. Moreover, since $f(\omega_k)$ converges to l , we also have $V_{\omega_k}(0) \rightarrow l$. We can guarantee convergence of $V_{\omega_k}(n)$ for $n > 0$ by passing to a subsequence because $R^{\mathbb{Z}}$ is compact. Let us denote this limit potential by V_l . As each V_{ω_k} lies in $\text{supp } \nu$, so does V_l ; moreover, $V(0) = l$ and $V_l(n) = V_{\omega_0}(n)$ for each $n < 0$. \square

Here is an illustration of this result and a strengthening of the derived consequence:

Corollary 5. *Suppose $\Omega = \mathbb{T}$, μ is normalized Lebesgue measure, and $T\omega = \omega + \alpha$ for some irrational α . If f has a single (non-removable) discontinuity at ω_0 , then for all $\omega \in [0, 1)$, the operator H_ω has no absolutely continuous spectrum.*

Proof. Let us say that l is a limiting value of f at ω_0 if there is a sequence $\{\omega_k\}$ in $\mathbb{T} \setminus \{\omega_0\}$ such that $\omega_k \rightarrow \omega_0$ and $f(\omega_k) \rightarrow l$. As f has a non-removable discontinuity at ω_0 , it has more than one limiting value at this point. Moreover, since f is continuous away from ω_0 , any limiting value is also an essential limit since we can choose each Ω_k to be a suitably small interval around ω_k .

This shows that f has an essential discontinuity at ω_0 . As the orbit of ω_0 never returns to this point, f is continuous at each point $T^n \omega_0$, $n \neq 0$. Therefore, Theorem 10 is applicable and shows that H_ω has no absolutely continuous spectrum for Lebesgue-almost every $\omega \in [0, 1)$.

It remains to show that the absolutely continuous spectrum of H_ω is empty for all $\omega \in \mathbb{R}/\mathbb{Z}$. We begin by fixing ω_1 such that H_{ω_1} has no absolutely continuous spectrum and such that the orbit of ω_1 does not meet ω_0 ; almost all ω_1 have these properties.

Given an arbitrary $\omega \in \mathbb{R}/\mathbb{Z}$, we may choose a sequence of integers $\{n_i\}$ so that $T^{n_i}(\omega) \rightarrow \omega_1$. As f is continuous on the orbit of ω_1 , the potentials associated to $T^{n_i}(\omega)$ converge pointwise to V_{ω_1} . By a result of Last and Simon [51], the absolutely continuous spectrum cannot shrink under pointwise approximation using translates of a single potential. Thus, the operator with potential V_ω cannot have absolutely continuous spectrum. This concludes the proof. \square

These results are particularly interesting in connection with Problem 2. Quasi-periodic potentials (as defined in this paper) are almost periodic if and only if f is continuous. Thus, proving the absence of absolutely continuous spectrum for quasi-periodic potentials with discontinuous f 's is a way of providing further support for the conjecture that $\Sigma_{ac} \neq \emptyset$ implies almost periodicity.

Let us now turn to the case of continuous sampling functions f . The proof of Theorem 10 certainly breaks down and it is not clear where some sort of non-determinism should come from in the quasi-periodic case, for example. Of course, absence of absolutely continuous spectrum does not hold for a general continuous f . Thus, the following result from [2] is somewhat surprising:

Theorem 11 (Avila-Damanik 2005). *Suppose Ω is a compact metric space, $T : \Omega \rightarrow \Omega$ a homeomorphism, and μ a non-atomic ergodic Borel probability measure. Then, there is a residual set of functions f in $C(\Omega)$ such that $\Sigma_{ac}(f) = \emptyset$.*

Recall that a subset of $C(\Omega)$ is called residual if it contains a countable intersection of dense open sets. A residual set is locally uncountable.

One would expect some absolutely continuous spectrum for weak perturbations with sufficiently nice potentials; especially in the one-frequency quasi-periodic case. More precisely, if f is nice enough, then $\Delta + \lambda f(n\alpha + \omega)$ should have some/purely absolutely continuous spectrum for $|\lambda|$ sufficiently small. It is known that real-analyticity is sufficiently “nice enough” [15] (when α is Diophantine), but it was expected that this assumption is much too strong and could possibly be replaced by mere continuity. The proof of Theorem 11 can easily be adapted to yield the following result, also contained in [2], which shows that continuity of the sampling function is not sufficient to ensure the existence of absolutely continuous spectrum for weakly coupled quasi-periodic potentials.

Theorem 12 (Avila-Damanik 2005). *Suppose Ω is a compact metric space, $T : \Omega \rightarrow \Omega$ a homeomorphism, and μ a non-atomic ergodic Borel probability measure. Then, there is a residual set of functions f in $C(\Omega)$ such that $\Sigma_{ac}(\lambda f) = \emptyset$ for almost every $\lambda > 0$.*

Proof. We only sketch the proofs of Theorems 11 and 12. The key technical issue is to establish that the maps

$$(31) \quad (L^1(\Omega) \cap B_r(L^\infty(\Omega)), \|\cdot\|_1) \rightarrow \mathbb{R}, \quad f \mapsto \mathcal{Z}(f)$$

and

$$(32) \quad (L^1(\Omega) \cap B_r(L^\infty(\Omega)), \|\cdot\|_1) \rightarrow \mathbb{R}, \quad f \mapsto \int_0^\Lambda \mathcal{Z}(\lambda f) d\lambda$$

are upper semi-continuous. Here, $\Lambda > 0$, $B_r(L^\infty(\Omega)) = \{f \in L^\infty(\Omega) : \|f\|_\infty < r\}$, and $\mathcal{Z}(f)$ denotes the set of energies for which the Lyapunov exponent associated with (Ω, T, μ, f) vanishes.

Upper semi-continuity of the map (31) can be shown using the fact that γ is harmonic in the upper half-plane and subharmonic on the real line; see [2] for details. Fatou's Lemma then implies upper semi-continuity of (32).

For $\delta > 0$, define

$$M_\delta = \{f \in C(\Omega) : \text{Leb}(\mathcal{Z}(f)) < \delta\}.$$

By the upper semi-continuity statement above, M_δ is open. By approximation with discontinuous functions and upper semi-continuity again, we see that M_δ is also dense.

It follows that

$$\{f \in C(\Omega) : \Sigma_{\text{ac}}(f) = \emptyset\} = \{f \in C(\Omega) : \text{Leb}(\mathcal{Z}(f)) = 0\} = \bigcap_{\delta > 0} M_\delta$$

is residual and Theorem 11 follows. Given upper semi-continuity of (32), the proof of Theorem 12 is analogous. \square

While continuous functions can be approximated in the C^0 norm by discontinuous functions, this does not work in the C^ε norm for any $\varepsilon > 0$. Thus, the proof just given does not extend to Hölder classes. It would be interesting to explore possible extensions of the results themselves; thus motivating the following problem.

Problem 7. *Prove or disprove statements like the ones in Theorems 11 and 12 for Hölder classes $C^\varepsilon(\Omega)$, $\varepsilon > 0$.*

6. UNIFORM LYAPUNOV EXPONENTS AND ZERO-MEASURE SPECTRUM

The Kotani result for potentials taking finitely many values, Theorem 9, is central to the study of one-dimensional quasi-crystal models. The main results in this area have been reviewed in [24, 25, 63]. In this section we will therefore focus on the recent progress and discuss why zero-measure spectrum is a consequence of Kotani theory when there is uniform convergence to the Lyapunov exponent.

One-dimensional quasi-crystals are typically modelled by sequences over a finite alphabet which are aperiodic but which have very strong long-range order properties. An important class of examples is given by one-frequency quasi-periodic potentials with step functions as sampling functions. That is, the potentials are of the form

$$(33) \quad V_\omega(n) = \lambda \sum_{m=1}^N \gamma_m \chi_{[a_{m-1}, a_m)}(n\alpha + \omega),$$

where $0 = a_0 < a_1 < \dots < a_N = 1$ is a partition of the unit circle, $\lambda, \gamma_1, \dots, \gamma_N$ are real numbers, α is irrational, and $\omega \in \mathbb{T}$. We obtain aperiodic potentials if $\lambda \neq 0$ and $\{\gamma_1, \dots, \gamma_N\}$ has cardinality at least two. We will assume these conditions throughout this section.

One of the properties that has been established for many quasi-crystal models is zero-measure spectrum. By general principles, this implies that the spectrum is a Cantor set because it cannot contain isolated points. Given the Kotani result, $\text{Leb}(\mathcal{Z}) = 0$, the natural way of proving this is via the identity $\Sigma = \mathcal{Z}$.

Theorem 13 (Damanik-Lenz 2006). *Suppose the potentials are of the form (33) and in addition all discontinuity points $\{a_m\} \subset \mathbb{T}$ are rational. Then, the Lyapunov exponent vanishes identically on the spectrum, that is, $\Sigma = \mathcal{Z}$. As a consequence, the spectrum is a Cantor set of zero Lebesgue measure.*

Proof. We only sketch the main ideas. More details can be found in [28, 29]. Denote

$$\mathcal{UH} = \left\{ E : \frac{1}{n} \log \|A_n^E(\omega)\| \rightarrow \gamma(E) > 0 \text{ uniformly in } \omega \right\}.$$

Then, by Lenz [53] (see also Johnson [40]), $\mathcal{UH} = \mathbb{C} \setminus \Sigma$. In particular,

$$\Sigma = \mathcal{Z} \cup \mathcal{N}\mathcal{UH},$$

where

$$\mathcal{NUH} = \{E : \gamma(E) > 0 \text{ and } E \notin \mathcal{UH}\}.$$

The result follows once $\mathcal{NUH} = \emptyset$ is established. Thus, given E with $\gamma(E) > 0$, we need to show that $\frac{1}{n} \log \|A_n^E(\omega)\| \rightarrow \gamma(E)$ uniformly in ω .

Uniform convergence along a special subsequence, $n_k \rightarrow \infty$, can be shown using results from Boshernitzan [13] and Lenz [52]. Namely, the assumption that all a_m are rational implies that there is a sequence of integers $n_k \rightarrow \infty$ such that for each k , all words of length n_k that occur in the potentials V_ω do so with comparable frequencies. That is, there is a uniform $C > 0$ such that for every k ,

$$(34) \quad \min_{|w|=n_k, w \text{ occurs}} \liminf_{J \rightarrow \infty} \frac{1}{J} \#\{j : 1 \leq j \leq J, V_\omega(j) \dots V_\omega(j+n_k-1) = w\} \geq \frac{C}{n_k}$$

uniformly in ω [13]. Using this result, one can then use ideas from [52] to show that $\frac{1}{n_k} \log \|A_{n_k}^E(\omega)\| \rightarrow \gamma(E)$ as $k \rightarrow \infty$, uniformly in ω .

Finally, the avalanche principle of Goldstein and Schlag [35] allows one to interpolate and prove the desired uniform convergence of $\frac{1}{n} \log \|A_n^E(\omega)\|$ to $\gamma(E)$ as $n \rightarrow \infty$. \square

The rationality assumption in Theorem 13 holds on a dense set of parameters, which makes it suitable for an approximation of a continuous sampling function by a sequence of step functions. Some consequences that may be drawn from this can be found in Bjerklöv et al. [12]. On the other hand, the assumption is certainly not necessary and more general results than the one presented here can be found in [29]. It would be nice if the assumption could be removed altogether:

Problem 8. *Prove zero-measure spectrum for all finite partitions of the circle, that is, remove the rationality assumption from Theorem 13.*

It should be mentioned, however, that the proof sketched above will not work in this generality. The approach is based on the Boshernitzan condition (34), and it was shown in [29] that this condition fails for certain parameter values.

From a mathematical point of view, the following problem is natural:

Problem 9. *Study multi-frequency analogs. That is, for a finite partition $\mathbb{T}^d = J_1 \cup \dots \cup J_N$ and the operators with potential*

$$V_\omega(n) = \lambda \sum_{m=1}^N \gamma_m \chi_{J_m}(n\alpha + \omega),$$

what is the measure of the spectrum and what is the spectral type?

These potentials are not directly motivated by quasi-crystal theory but they form an interesting class that may again prove useful in the understanding of the phenomena that arise for continuous sampling functions. Moreover, very little is understood about the associated operators apart from the general Kotani result, which says that there is never any absolutely continuous spectrum. It is unclear, however, whether there can be any point spectrum, for example.

7. PURELY AC SPECTRUM FOR THE SUBCRITICAL AMO

In this final section, we describe the (to the best of our knowledge) first application of Corollary 1. In his 1997 paper, Kotani writes that at the time Kotani

theory was developed, “it was not clear whether we could know the pure absolute continuity ... only from the IDS, but this corollary has answered this question affirmatively.”

The application we will present involves the almost Mathieu operator

$$[H_\omega \psi](n) = \psi(n+1) + \psi(n-1) + 2\lambda \cos(2\pi(n\alpha + \omega))\psi(n),$$

that is, the Schrödinger operator with one-frequency quasi-periodic potential associated with the sampling function $f(\omega) = 2\lambda \cos(2\pi\omega)$. This operator is known to exhibit a metal-insulator transition at $|\lambda| = 1$, that is, for almost every α , the almost sure spectral type is purely absolutely continuous for $|\lambda| < 1$, purely singular continuous for $|\lambda| = 1$, and pure point (with exponentially decaying eigenfunctions) for $|\lambda| > 1$. See Jitomirskaya [38] for this result and its history. For $|\lambda| > 1$, one indeed has to exclude a zero-measure set of frequencies α since it was shown by Avron and Simon, using Gordon’s Lemma, that for Liouville α , there are no eigenvalues [8, 9]. For $|\lambda| \leq 1$, it has long been expected that the results above extend to all irrational frequencies. For $|\lambda| = 1$, the issue was resolved by Avila and Krikorian [6]. The question of what happens for $|\lambda| < 1$ was addressed by Problem 6 in [60].

Since Bourgain and Jitomirskaya showed in [16] that the Lyapunov exponent associated with the almost Mathieu operator obeys $\gamma(E) = \max\{0, \log |\lambda|\}$ for every $E \in \Sigma$ for all irrational frequencies α , the problem reduces to a study of the integrated density of states, that is, to a proof of its absolute continuity. The following result was shown in [3] and it completely settles this regularity issue for the integrated density of states.

Theorem 14 (Avila-Damanik 2006). *The integrated density of states of the almost Mathieu operator is absolutely continuous if and only if $|\lambda| \neq 1$.*

Combining this result with the one from [16] just quoted along with Corollary 1, we obtain almost surely purely absolutely continuous spectrum for the subcritical (i.e., $|\lambda| < 1$) almost Mathieu operator:

Corollary 6. *If $|\lambda| < 1$, then $\Sigma_{\text{sing}} = \emptyset$.*

Subsequently, Avila even extended this result and proved purely absolutely continuous spectrum for $|\lambda| < 1$, α irrational, and every ω [1].

Problem 10. *Extend the results of this section to more general $f \in L^\infty(\mathbb{T})$; for example, real-analytic f ’s.*

It was shown by Bourgain and Jitomirskaya that for Diophantine frequency α and analytic f , $\Sigma_{\text{sing}}(\lambda f) = \emptyset$ for λ sufficiently small [15]. An extension of this result to all ω is contained in [5]. The proofs probably break down for Liouville frequencies (for this, one needs to extend the method based on Gordon’s Lemma to matrices that are not banded, but which do have exponential off-diagonal decay). Thus, it seems natural to attack the problem for Liouville frequencies in the same way as above.

REFERENCES

- [1] A. Avila, Absolutely continuous spectrum for the almost Mathieu operator with subcritical coupling, in preparation
- [2] A. Avila and D. Damanik, Generic singular spectrum for ergodic Schrödinger operators, *Duke Math. J.* **130** (2005), 393–400

- [3] A. Avila and D. Damanik, Absolute continuity of the integrated density of states for the almost Mathieu operator with non-critical coupling, preprint (2006)
- [4] A. Avila and D. Damanik, in preparation
- [5] A. Avila and S. Jitomirskaya, Almost localization and almost reducibility, in preparation
- [6] A. Avila and R. Krikorian, Reducibility or non-uniform hyperbolicity for quasiperiodic Schrödinger cocycles, to appear in *Ann. of Math.*
- [7] J. Avron and B. Simon, Almost periodic Schrödinger operators. I. Limit periodic potentials, *Commun. Math. Phys.* **82** (1981), 101–120
- [8] J. Avron and B. Simon, Singular continuous spectrum for a class of almost periodic Jacobi matrices, *Bull. Amer. Math. Soc.* **6** (1982), 81–85
- [9] J. Avron and B. Simon, Almost periodic Schrödinger operators. II. The integrated density of states, *Duke Math. J.* **50** (1983), 369–391
- [10] J. Bochi, Genericity of zero Lyapunov exponents, *Ergod. Th. & Dynam. Sys.* **22** (2002), 1667–1696
- [11] J. Bochi and M. Viana, The Lyapunov exponents of generic volume-preserving and symplectic maps, *Ann. of Math.* **161** (2005), 1–63
- [12] K. Bjerklov, D. Damanik, and R. Johnson, Lyapunov exponents of continuous Schrödinger cocycles over irrational rotations, preprint (2005)
- [13] M. Boshernitzan, Rank two interval exchange transformations, *Ergod. Th. & Dynam. Sys.* **8** (1988), 379–394
- [14] J. Bourgain, Positivity and continuity of the Lyapunov exponent for shifts on \mathbb{T}^d with arbitrary frequency vector and real analytic potential, *J. Anal. Math.* **96** (2005), 313–355
- [15] J. Bourgain and S. Jitomirskaya, Absolutely continuous spectrum for 1D quasiperiodic operators, *Invent. Math.* **148** (2002), 453–463
- [16] J. Bourgain and S. Jitomirskaya, Continuity of the Lyapunov exponent for quasiperiodic operators with analytic potential, *J. Statist. Phys.* **108** (2002), 1203–1218
- [17] J. Bourgain and W. Schlag, Anderson localization for Schrödinger operators on \mathbb{Z} with strongly mixing potentials, *Commun. Math. Phys.* **215** (2000), 143–175
- [18] D. Buschmann, A proof of the Ishii-Pastur theorem by the method of subordination, *Univ. Jägel. Acta Math.* **34** (1997), 29–34
- [19] A. Casher and J. Lebowitz, Heat flow in regular and disordered harmonic chains, *J. Math. Phys.* **12** (1971), 1701–1711
- [20] V. Chulaevsky, An inverse spectral problem for limit-periodic Schrödinger operators, *Functional Anal. Appl.* **18** (1984), 220–233
- [21] V. Chulaevsky and S. Molchanov, The structure of a spectrum of the lacunary-limit-periodic Schrödinger operator, *Functional Anal. Appl.* **18** (1984), 343–344
- [22] V. Chulaevsky and T. Spencer, Positive Lyapunov exponents for a class of deterministic potentials, *Commun. Math. Phys.* **168** (1995), 455–466
- [23] H. Cycon, R. Froese, W. Kirsch, and B. Simon, *Schrödinger Operators with Application to Quantum Mechanics and Global Geometry*, Texts and Monographs in Physics, Springer-Verlag, Berlin, 1987
- [24] D. Damanik, Gordon-type arguments in the spectral theory of one-dimensional quasicrystals, in *Directions in Mathematical Quasicrystals*, CRM Monograph Series **13**, American Mathematical Society, Providence (2000), 277–305
- [25] D. Damanik, Strictly ergodic subshifts and associated operators, to appear in *Simon Festschrift*, Proceedings of Symposia in Pure Mathematics, American Mathematical Society, Providence
- [26] D. Damanik and R. Killip, Ergodic potentials with a discontinuous sampling function are non-deterministic, *Math. Res. Lett.* **12** (2005), 187–192
- [27] D. Damanik and R. Killip, Almost everywhere positivity of the Lyapunov exponent for the doubling map, *Commun. Math. Phys.* **257** (2005), 287–290
- [28] D. Damanik and D. Lenz, A criterion of Boshernitzan and uniform convergence in the multiplicative ergodic theorem, *Duke Math. J.* **133** (2006), 95–123
- [29] D. Damanik and D. Lenz, Zero-measure Cantor spectrum for Schrödinger operators with low-complexity potentials, *J. Math. Pures Appl.* **85** (2006), 671–686
- [30] P. Deift and B. Simon, Almost periodic Schrödinger operators. III. The absolutely continuous spectrum in one dimension, *Commun. Math. Phys.* **90** (1983), 389–411

- [31] J. Geronimo, Polynomials orthogonal on the unit circle with random recurrence coefficients, *Methods of Approximation Theory in Complex Analysis and Mathematical Physics* (Leningrad, 1991), 43–61, Lecture Notes in Math., 1550, Springer, Berlin, 1993
- [32] J. Geronimo and A. Teplyaev, A difference equation arising from the trigonometric moment problem having random reflection coefficients—an operator-theoretic approach, *J. Funct. Anal.* **123** (1994), 12–45
- [33] D. Gilbert, On subordinacy and analysis of the spectrum of Schrödinger operators with two singular endpoints, *Proc. Roy. Soc. Edinburgh A* **112** (1989), 213–229
- [34] D. Gilbert and D. Pearson, On subordinacy and analysis of the spectrum of one-dimensional Schrödinger operators, *J. Math. Anal. Appl.* **128** (1987), 30–56
- [35] M. Goldstein and W. Schlag, Hölder continuity of the integrated density of states for quasi-periodic Schrödinger equations and averages of shifts of subharmonic functions, *Ann. of Math.* **154** (2001), 155–203
- [36] M. Herman, Une méthode pour minorer les exposants de Lyapunov et quelques exemples montrant le caractère local d’un théorème d’Arnold et de Moser sur le tore de dimension 2, *Comment. Math. Helv.* **58** (1983), 4453–502
- [37] K. Ishii, Localization of eigenstates and transport phenomena in the one dimensional disordered system, *Supp. Theor. Phys.* **53** (1973), 77–138
- [38] S. Jitomirskaya, Metal-insulator transition for the almost Mathieu operator, *Ann. of Math.* **150** (1999), 1159–1175
- [39] S. Jitomirskaya, Ergodic Schrödinger operators (on one foot), to appear in *Simon Festschrift*, Proceedings of Symposia in Pure Mathematics, American Mathematical Society, Providence
- [40] R. Johnson, Exponential dichotomy, rotation number, and linear differential operators with bounded coefficients, *J. Differential Equations* **61** (1986), 54–78
- [41] W. Kirsch, On a class of random Schrödinger operators, *Adv. in Appl. Math.* **6** (1985), 177–187
- [42] W. Kirsch, S. Kotani and B. Simon, Absence of absolutely continuous spectrum for some one-dimensional random but deterministic Schrödinger operators, *Ann. Inst. H. Poincaré Phys. Théor.* **42** (1985), 383–406
- [43] S. Kotani, Ljapunov indices determine absolutely continuous spectra of stationary random one-dimensional Schrödinger operators, *Stochastic Analysis* (Katata/Kyoto, 1982), 225–247, North-Holland Math. Library **32**, North-Holland, Amsterdam, 1984
- [44] S. Kotani, Support theorems for random Schrödinger operators, *Commun. Math. Phys.* **97** (1985), 443–452
- [45] S. Kotani, One-dimensional random Schrödinger operators and Herglotz functions, *Probabilistic methods in mathematical physics* (Katata/Kyoto, 1985), 219–250, Academic Press, Boston, MA, 1987
- [46] S. Kotani, Jacobi matrices with random potentials taking finitely many values, *Rev. Math. Phys.* **1** (1989), 129–133
- [47] S. Kotani, Generalized Floquet theory for stationary Schrödinger operators in one dimension, *Chaos Solitons Fractals* **8** (1997), 1817–1854
- [48] S. Kotani and M. Krishna, Almost periodicity of some random potentials, *J. Funct. Anal.* **78** (1988), 390–405
- [49] S. Kotani and B. Simon, Stochastic Schrödinger operators and Jacobi matrices on the strip, *Commun. Math. Phys.* **119** (1988), 403–429
- [50] H. Kunz and B. Souillard, Sur le spectre des opérateurs aux différences finies aléatoires, *Commun. Math. Phys.* **78** (1980/81), 201–246
- [51] Y. Last and B. Simon, Eigenfunctions, transfer matrices, and absolutely continuous spectrum of one-dimensional Schrödinger operators, *Invent. Math.* **135** (1999), 329–367
- [52] D. Lenz, Uniform ergodic theorems on subshifts over a finite alphabet, *Ergod. Th. & Dynam. Sys.* **22** (2002), 245–255
- [53] D. Lenz, Singular continuous spectrum of Lebesgue measure zero for one-dimensional quasicrystals, *Commun. Math. Phys.* **227** (2002), 119–130
- [54] N. Minami, An extension of Kotani’s theorem to random generalized Sturm-Liouville operators, *Commun. Math. Phys.* **103** (1986), 387–402
- [55] V. Oseledec, A multiplicative ergodic theorem. Characteristic Ljapunov exponents of dynamical systems (Russian), *Trudy Moskov. Mat. Obšč.* **19** (1968), 179–210; Translation in *Trans. Mosc. Math. Soc.* **19** (1968), 197–231

- [56] L. Pastur, Spectral properties of disordered systems in the one-body approximation, *Commun. Math. Phys.* **75** (1980), 179–196
- [57] L. Pastur and A. Figotin, *Spectra of Random and Almost-Periodic Operators*, Grundlehren der Mathematischen Wissenschaften **297**, Springer-Verlag, Berlin (1992)
- [58] L. Pastur and V. Tkachenko, Spectral theory of a class of one-dimensional Schrödinger operators with limit-periodic potentials (Russian), *Trudy Moskov. Mat. Obšč.* **51** (1988), 114–168, 258; Translation in *Trans. Moscow Math. Soc.* (1989), 115–166
- [59] B. Simon, Kotani theory for one-dimensional stochastic Jacobi matrices, *Commun. Math. Phys.* **89** (1983), 227–234
- [60] B. Simon, Schrödinger operators in the twenty-first century, *Mathematical Physics 2000*, 283–288, Imp. Coll. Press, London, 2000
- [61] B. Simon, *Trace Ideals and Their Applications*, Second edition, Mathematical Surveys and Monographs **120**, American Mathematical Society, Providence, RI, 2005
- [62] E. Sorets and T. Spencer, Positive Lyapunov exponents for Schrödinger operators with quasi-periodic potentials, *Commun. Math. Phys.* **142** (1991), 543–566
- [63] A. Sütő, Schrödinger difference equation with deterministic ergodic potentials, in *Beyond Quasicrystals* (Les Houches, 1994), Springer, Berlin (1995), 481–549
- [64] L.-S. Young, Some open sets of nonuniformly hyperbolic cocycles, *Ergod. Th. & Dynam. Sys.* **13** (1993), 409–415

MATHEMATICS 253–37, CALIFORNIA INSTITUTE OF TECHNOLOGY, PASADENA, CA 91125, USA
E-mail address: `damanik@caltech.edu`
URL: <http://www.math.caltech.edu/people/damanik.html>