

# MODULAR FORMS AND CALABI-YAU VARIETIES

KAPIL PARANJAPE<sup>1</sup> AND DINAKAR RAMAKRISHNAN<sup>2</sup>

## INTRODUCTION

Let  $f(z) = \sum_{n=1}^{\infty} a_n q^n$  be a holomorphic newform of weight  $k \geq 2$  relative to  $\Gamma_1(N)$  acting on the upper half plane  $\mathcal{H}$ . Suppose the coefficients  $a_n$  are all rational. When  $k = 2$ , a celebrated theorem of Shimura asserts that there corresponds an elliptic curve  $E$  over  $\mathbb{Q}$  such that for all primes  $p \nmid N$ ,  $a_p = p + 1 - |E(\mathbb{F}_p)|$ . Equivalently, there is, for every prime  $\ell$ , an  $\ell$ -adic representation  $\rho_\ell$  of the absolute Galois group  $\mathfrak{G}_{\mathbb{Q}}$  of  $\mathbb{Q}$ , given by its action on the  $\ell$ -adic Tate module of  $E$ , such that  $a_p$  is, for any  $p \nmid \ell N$ , the trace of the Frobenius  $Fr_p$  at  $p$  on  $\rho_\ell$ . Denote by  $t = t(N)$  the order of the torsion subgroup of  $\Gamma_1(N)$ .

The primary aim of this article is to provide some positive evidence for the expectation of Mazur and van Stratten that for every  $k \geq 2$ , any rational newform  $f$  of weight  $k$  and level  $N$  should have an associated Calabi-Yau variety  $X/\mathbb{Q}$  of dimension  $k - 1$  such that

- (Ai) The  $\{(k - 1, 0), (0, k - 1)\}$ -piece of  $H^{k-1}(X)$  splits off as a sub-motive  $M_f$  over  $\mathbb{Q}$ ,
- (Aii)  $a_p = \text{tr}(Fr_p | M_{f,\ell})$ , for all  $p$  not dividing  $\ell t N$ , and
- (Aiii)  $\det(M_{f,\ell}) = \chi_\ell^{k-1}$ ,

where  $\chi_\ell$  is the  $\ell$ -adic cyclotomic character.

In fact, we even hope that in addition the following holds:

- (Aiv)  $X$  admits an involution  $\tau$  which acts by  $-1$  on  $H^0(X, \Omega^{k-1})$ .

The hope is that the quotient  $X/\tau$  is a rational variety.

This typically holds in our examples below. This extra structure is natural to want and is needed for understanding twists. It obviously exists for  $k = 2$ , in which case  $X$  is (by Shimura) an elliptic curve over  $\mathbb{Q}$ , given by an equation  $y^2 = f(x)$ , and the involution  $\tau$  sends  $(x, y)$  to  $(-x, y)$ , thus acting by  $-1$  on the holomorphic differential  $\omega = dx/y$  which spans  $H^0(X, \Omega^1)$ . (Of course  $X/\tau$  is in this case  $\mathbb{P}^1$ .)

---

<sup>1</sup> Partly supported by the the JCBose Fellowship of the DST (IISERM:10-DST-JCB-F.3) and IMSC, Chennai.

<sup>2</sup> Partly supported by the NSF grants DMS-0701089 and DMS-1001916.

To fix ideas, one could think of a motive as a semisimple motive  $M$  relative to absolute Hodge cycles ([DMOS]), with realizations  $(M_B, M_{\text{dR}}, M_\ell)$ . Since every Calabi-Yau manifold of dimension 1 is an elliptic curve, (Ai) through (Aiv) provide a natural extension of what one has for  $k = 2$ . Of course for  $k > 2$ , one knows by Deligne ([Del]) that there is an irreducible, 2-dimensional  $\ell$ -adic representation  $\rho_\ell$  so that (Aii), (Aiii) hold with  $M_{f,\ell}$  replaced by  $\rho_\ell$ .

In the first part of this article we will focus on the forms  $f$  of even weight case for small levels, before moving on in the second part to formulate an analogue for regular selfdual cusp forms on  $\text{GL}(n)/\mathbb{Q}$  and see how the framework is compatible with the principle of functoriality.

When the weight is odd, the  $\mathbb{Q}$ -rationality forces  $f$  to be of CM type, and for weight 3, we refer to the paper of Elkies and Schuett ([ES]) for a beautiful result.

For non-CM newforms  $f$  of weight  $k > 2$  with  $\mathbb{Q}$ -coefficients, we in fact hope for more, namely that the cohomology ring of  $X$  will be spanned by  $M$  and the Hodge/Tate classes of various degrees; in particular,  $X$  should be rigid in this case.

It is a difficult problem in dimensions  $> 3$  to find smooth models of varieties with trivial canonical bundles, and for this reason we formulate our questions for such varieties with mild singularities. By a *Calabi-Yau variety* over a field  $k$ , we will mean an  $n$ -dimensional projective variety  $X/k$  on which the canonical bundle  $\mathcal{K}_X$  is defined such that

- (CY1)  $\mathcal{K}_X$  is trivial; and
- (CY2)  $H^m(X, \mathcal{O}_X) = 0$  for all (strictly) positive  $m < n$ .

More precisely, we will want such an  $X$  to be normal and Cohen-Macaulay, so that the dualizing sheaf  $\mathcal{K}_X$  is defined, with the singular locus in codimension at least 2, so that  $\mathcal{K}_X$  defines a Weil divisor; finally,  $X$  should be Gorenstein, so that  $\mathcal{K}_X$  will represent a Cartier divisor. Ideally we would like to singular locus  $X_{\text{sing}}$  to be of dimension  $\leq \lfloor \frac{n-1}{2} \rfloor$ .

In addition to these properties, we would ideally also like  $X$  to be realized as a double cover  $\pi : X \rightarrow Y$ , where  $Y$  is a projective smooth rational variety with negative ample canonical divisor. (Such an  $X$  will be automatically Gorenstein so that  $\mathcal{K}_X$  is defined, and  $\pi_*(\mathcal{O}_X) = \mathcal{O}_Y \oplus L$ , for a line bundle  $L$  on  $Y$  with  $L^2$  giving the branch locus; moreover,  $\pi_*(\mathcal{K}_X) = \mathcal{K}_Y \oplus \mathcal{K}_Y \otimes L^{-1}$ , forcing  $L = \mathcal{K}_Y$ .)

Here is our first result:

**Theorem 1** *Fix  $\Gamma = \Gamma_1(N)$ ,  $N \leq 5$ . Let  $k$  be the first even weight s.t.  $\dim(S_k(\Gamma)) = 1$ . Then  $\exists$  a Calabi-Yau variety  $V(f)/\mathbb{Q}$  with an*

involution  $\tau$  associated to the new generator  $f$  of  $S_k(\Gamma)$  satisfying (Ai) and (Aiv). In fact, when  $N \leq 5$ ,  $V = V(f)$  is birational over  $\mathbb{Q}$  to the Kuga-Sato variety  $\tilde{\mathcal{E}}_N^{(k-2)}$ .

In particular, this result applies to the Delta function  $\Delta(z) = \sum_{n=1}^{\infty} \tau(n)q^n = \prod_{m \geq 1} (1 - q^m)^{24}$ , for  $N = 1, k = 12$ . By the Kuga variety, we mean a suitable compactification (see section 1) of the fibre product  $E_N^{k-2}$  of the universal elliptic curve  $E_N$  over the model over  $\mathbb{Q}$  of the modular curve  $\Gamma_1(N) \backslash \mathcal{H}$ .

It is well known that  $S_k(\Gamma_1(N))$  is **one-dimensional** when  $(N, k)$  equals **(1, 12)**, (1, 16), **(2, 8)**, (2, 10), **(3, 6)**, (3, 8), **(4, 6)**, **(5, 4)**, (5, 6), **(6, 4)**, (7, 4). (The ones in bold are the cases to which the Theorem applies.) For example, for the case **(2, 8)**, the generator is

$$f(z) = q - 8q^2 + 12q^3 - 210q^4 + 1016q^5 + \dots$$

Recall that a newform  $f(z) = \sum_{n=1}^{\infty} a_n q^n$  is of *CM-type* iff there is an odd, quadratic Dirichlet character  $\delta$  such that  $a_p = a_p \delta(p)$  for almost all primes  $p$ . Equivalently, if  $K$  is the imaginary quadratic field cut out by  $\delta$ ,  $a_p = 0$  for all  $p$  which are inert in  $K$ .

As a first step, we may ask for a *potential statement*, i.e., the association, over a finite extension  $k$  of  $\mathbb{Q}$ , of a Calabi-Yau variety  $V/k$  to a newform  $f$  with  $\mathbb{Q}$ -coefficients. Here we state a modest result in this direction, already known in different ways, just to show that it fits into our framework:

**Proposition 2** Let  $f$  be a newform of weight  $k \geq 3$  of CM type with rational coefficients. Then  $\exists$  a Calabi-Yau  $(k - 1)$ -fold  $X$  defined over a number field  $F$  such that (Ai), (Aii) and (Aiii) hold over  $F$ . This  $X$  arises as a Kummer variety associated to an elliptic curve  $E$  with complex multiplication. When  $k \leq 4$ ,  $X$  can be taken to be a smooth model.

Again, when  $k = 3$ , there is a much more precise and satisfactory result over  $\mathbb{Q}$  in the work of Elkies and Schuett [ES].

In the converse direction, if  $M$  is a simple motive over  $\mathbb{Q}$  of rank 2, with coefficients in  $\mathbb{Q}$ , of Hodge type  $\{(w, 0), (0, w)\}$  with  $w > 0$ , then the general philosophy of Langlands, and also a conjecture of Serre, predicts that  $M$  should be modular and be associated to a newform  $f$  of weight  $w + 1$  with rational coefficients. This is part of a very general phenomenon, and applies to motives occurring in the cohomology smooth projective varieties over  $\mathbb{Q}$ . In any case, it applies in particular to Calabi-Yau threefolds over  $\mathbb{Q}$  whose  $\{(3, 0), (0, 3)\}$ -part splits off as

a submotive. In this context, there have been a number of beautiful results, some of which have been described in the monographs [YL] and [YYL]. They are entirely consistent with what we are trying to do in the opposite direction, and also provide supporting examples. It should perhaps be remarked that these results (relating to the modularity of rigid Calabi-Yau threefolds) can now be deduced *en masse* from the proof of Serre's conjecture due to Khare and Wintenberger ([KW]), with a key input from Kisin ([Kis]).

Let us now move to a more general situation. Fix any positive integer  $n$  and suppose that  $f$  is a (new) Hecke eigen-cuspform on the symmetric space

$$\mathcal{D}_n := \mathrm{SL}(n, \mathbb{R})/\mathrm{SO}(n),$$

relative to a congruence subgroup  $\Gamma$  of  $\mathrm{SL}(n, \mathbb{Z})$ , which is *algebraic* and *regular*. For  $n = 2$ ,  $f$  is algebraic and regular iff it is holomorphic of weight  $\geq 2$ . In general, one considers the cuspidal automorphic representation  $\pi$  of  $\mathrm{GL}(n, \mathbb{A})$  which is generated by  $f$ , and by Langlands the archimedean component  $\pi_\infty$  corresponds to an  $n$ -dimensional representation  $\sigma_\infty$  of the real Weil group  $W_{\mathbb{R}}$ , which contains  $\mathbb{C}^*$  as a subgroup of index 2. One says ([Clo1]) that  $\pi$  is algebraic if the restriction of  $\sigma_\infty$  to  $\mathbb{C}^*$  is a sum of characters  $\chi_j$  of the form  $z \rightarrow z^{p_j} \bar{z}^{q_j}$ , with  $p_j, q_j \in \mathbb{Z}$ , and it is regular iff  $\chi_i \neq \chi_j$  when  $i \neq j$ . Such an  $f$  contributes to the *cuspidal cohomology*  $H_{\mathrm{cusp}}^*(\Gamma \backslash \mathcal{D}_n, V)$  relative to a *local coefficient system*  $V$  in a specific degree  $w = w(f)$ . Moreover,  $f$  is rational over a number field  $\mathbb{Q}(f)$ , defined by the Hecke action on cohomology, which preserves the cuspidal part (*loc. cit.*). There is conjecturally a motive  $M(f)$  over  $\mathbb{Q}$  of rank  $n$ , with coefficients in  $\mathbb{Q}(f)$ , and weight  $w$ . By Clozel [Clo2],  $M_\ell(f)$  exists for suitable  $f$  which are in addition *essentially selfdual*.

Now let  $f$  be an algebraic, regular, essentially selfdual newform of weight  $w$  relative to  $\Gamma \subset \mathrm{SL}(n, \mathbb{Z})$ , with  $L$ -function  $L(s, f) = \prod_p L_p(s, f)$ , such that  $\mathbb{Q}(f) = \mathbb{Q}$ . Then our question is if there exists a Calabi-Yau variety  $X/\mathbb{Q}$  of dimension  $w$  with an involution  $\tau$  such that

- (Ci) There is a submotive  $M(f)$  of  $H^w(X)$  of rank  $n$  such that  $M(f)^{(w,0)} = H^{w,0}(X)$ ,
- (Cii)  $L_p(s, f) = L_p(s, M_\ell(f))$  for almost all  $p$ , and
- (Ciii) the quotient of  $X$  by  $\tau$  is a rational variety.

where  $L_p(s, M_\ell(f))$  equals, at any prime  $p \neq \ell$  where the  $\ell$ -adic realization  $M_\ell(f)$  is unramified,  $\det(I - Fr_p p^{-s} | M_\ell(f))^{-1}$ . Again we would like to be able to find an  $X$  having good reduction outside the primes

dividing  $tN$ , where  $n$  is the level of  $f$  and  $t$  the order of torsion in  $\Gamma$ , such that (Cii) holds for any such  $p \neq \ell$ .

Thanks to the *principle of functoriality*, one should be able to obtain a certain class of  $\mathbb{Q}$ -rational, regular, algebraic, essentially selfdual newforms  $f$  by transferring forms on (the symmetric domains of) smaller reductive  $\mathbb{Q}$ -subgroups  $G$  of  $\mathrm{GL}(n)$ . The simplest instance of this phenomenon is given by the symmetric powers  $\mathrm{sym}^m(g)$  of classical  $\mathbb{Q}$ -rational, non-CM newforms  $g$  of weight  $k$ . One knows by Kim and Shahidi ([KS], [Kim]) that for  $m \leq 4$ ,  $f = \mathrm{sym}^m(g)$  is a cusp form on  $\mathrm{GL}(m+1)$ . (Recently, this has been extended to  $m = 5$  in the works of Clozel-Thorne and Dieulefait.) Here is our third result:

**Theorem 3** *Let  $g$  be a non-CM, elliptic modular newform of weight 2, level  $N$  and trivial character, whose coefficients  $a_n$  lie in  $\mathbb{Q}$ . Then for any  $m > 0$ , there is a Calabi-Yau variety  $X_m$  with an involution  $\tau$  over  $\mathbb{Q}$  of dimension  $m$  associated to  $(g, \mathrm{sym}^m)$  such that (Ci), (Cii), (Ciii) hold relative to  $M_\ell = \mathrm{sym}^m(\rho_\ell(g))$ . Moreover, for  $m \leq 3$ ,  $X_m$  can be taken to be non-singular, with good reduction outside  $m!N$ .*

Here  $X_2$  is just the familiar Kummer surface attached to  $E \times E$ , where  $E = X_1$  is the elliptic curve/ $\mathbb{Q}$  defined by  $g$ . But the case  $m = 3$  is interesting, especially since it is not rigid, thanks to the Hodge type being  $\{(3,0), (2,1), (1,2), (0,3)\}$ , with each Hodge piece being one-dimensional. In fact, in that case,  $\mathrm{sym}^3(g)$  corresponds (by [RS]) to a holomorphic Siegel modular cusp form  $F$  of genus 2 and (Siegel) weight 3. Such an  $F$  contributes to the cohomology in degree 3 of the Siegel modular threefold  $V$  of level  $N^3$ . Since the geometric genus of  $V$  is typically  $> 1$ , it cannot be Calabi-Yau. However, there should be, as predicted by the Hodge and Tate conjectures, an algebraic correspondence between  $X_3$  and  $V$  (for any  $N$ ).

One also knows (cf. [Ram]) that given two non-CM newforms  $g, h$  of weights  $k, r \geq 2$  respectively, then there is an algebraic automorphic form  $f = g \boxtimes h$  on  $\mathrm{GL}(4)/\mathbb{Q}$ , which will be cuspidal and regular if  $k \neq r$ . If  $g, h$  are  $\mathbb{Q}$ -rational, then so is  $f$ . Moreover,  $f$  is essentially selfdual because  $g$  and  $h$  are.

**Theorem 4** *Let  $g, h$  be  $\mathbb{Q}$ -rational, non-CM newforms as above of respective weights  $k, r > 1$ , with  $k \neq r$ . Suppose we have Calabi-Yau varieties with involutions  $(X(g), \tau_g), (X(h), \tau_h)$  over  $\mathbb{Q}$  attached to  $g, h$  respectively, satisfying (Ai) through A(iv). Put  $f = g \boxtimes h$ , so that  $w(f) = (k-1)(r-1)$ . Then there is a Calabi-Yau variety with*

*involution  $(X(f), \tau_f)$  over  $\mathbb{Q}$  of dimension  $w(f)$  such that (Ci) through (Ciii) hold.*

The point is that the product  $Z := X(g) \times X(h)$  has the desired submotive in degree  $w$ , but it has global holomorphic  $m$ -forms for  $m = k - 1$  and  $m = r - 1$ . We exhibit an involution  $\tau$  on  $Z$  such that when we take the quotient by  $\tau$ , these forms get killed and we get a Calabi-Yau variety with reasonable singularities. To get unconditional examples of this Theorem, take  $k = 2$  and choose  $h$  to be one of the examples of Theorem A of weight  $r > 2$ . To be specific, we may take  $g$  to be the newform of weight 2 and level 11, and  $h$  to be the newform of weight 4 and level 5, in which case  $f = g \boxtimes h$  has level  $55^2$ , and  $X(f)$  is a Calabi-Yau fourfold.

In sum, it is a natural question, given Shimura's work on for of weight 2, if there are Calabi-Yau varieties with an involution associated to forms of higher weight with rational coefficients, and a preliminary version of this circle of questions was raised by the first author in a talk at the Borel memorial conference at Zhejiang University in Hangzhou, China, in 2004, and quite appropriately, Dick Gross, who was in the audience, cautioned against hoping for too much without sufficient evidence. Over the past years, there has been some positive evidence, though small, and even if there is no  $V$  in general, especially for non-CM forms  $f$  of even weight, the examples where one has nice Calabi-Yau varieties  $V$  enriches them considerably, and one of our aims is to understand the Hecke eigenvalues  $a_p$  in such cases a bit better in terms of counting points of  $V \bmod p$ . Since then we have learnt from [ES] (and Noriko Yui) that the question of existence of a Calabi-Yau variety  $V$  associated to  $f$  were earlier raised by Mazur and van Straten. What we truly hope for is that in addition,  $V$  will be equipped with an involution  $\tau$  acting by  $-1$  on the unique global holomorphic form of maximal degree, so one can form products, etc., and also deal with quadratic twists. We have some interesting examples in the 3-dimensional case (where  $k = 4$ ), and a sequel to this paper will also contain a discussion of these matters, as well as a way to get nicer models in certain higher dimensional examples.

The Ramanujan coefficients  $\tau(p)$  of the Delta function have been a source of much research. Our own work was originally motivated by the desire to express them in terms of the zeta function of a Calabi-Yau variety, though our path has diverged somewhat. In a different direction, a very interesting monograph of Edixhoven, et al ([?]) yields a deterministic algorithm for computing  $\tau(p)$  with expected running time

which is polynomial in  $\log p$ . They do this by relating the associated Galois representation mod  $\ell$  to the geometry of certain effective divisors on the modular curve  $X_1(5\ell)$ .

We thank all the people who have shown interest in this work, and a special thanks must go to Matthias Schuett who made a number of useful comments on the earlier version (2008). One of us (K.P.) would like to thank the JC Bose Fellowship of the DST (IISERM:10-DST-JCB-F.3) as well as Caltech and the IMSC, Chennai, where some of the work was done, and the second author (D.R.) would like to thank the NSF for continued support through the grants DMS-0701089 and DMS-1001916, and also IMSC, Chennai, for having him visit at various times.

## 1. AN INTUITIVE PICTURE OF THE GEOMETRIC CONSTRUCTION

In this section we will give an idea behind our construction of varieties with trivial canonical bundles arising as birational models of elliptic modular varieties  $V$  with non-positive canonical bundles. (S.T. Yau has informed us that he earlier had a similar construction, albeit in a different context.) We will use the intuitive language of divisors and linear systems. The content here will not be used in the succeeding sections, where we will use sheaves and do everything precisely in the different cases at hand.

Recall that  $V$  arise as fibre products of the universal family of elliptic curves  $E$  with additional structures over the modular curve associated to a congruence subgroup  $\Gamma$  of  $\mathrm{SL}(2, \mathbb{Z})$ . When  $\Gamma$  is  $\Gamma_1(N)$  (resp.  $\Gamma_0(N)$ ), the additional structure is a point (resp. subgroup) of order  $N$ . We have a slight preference for  $\Gamma_1(N)$  over  $\Gamma_0(N)$ , because in the latter case, one gets, due to the existence of  $-I$ , only a coarse moduli space.

Suppose we want to parametrize triples of the form  $(E, S; R)$  where  $E$  is a curve of genus 1,  $S$  a finite set of points on  $E$ , and  $R$  a finite set of linear equivalence relations on the points  $S$ . Suppose also that there is a surface  $X$ , and a linear system  $P$  of divisors linear equivalent to  $-K_X$ , where  $K_X$  is the canonical divisor of  $X$ . Assume that for a “general” datum  $(E, S; R)$  we have a uniquely determined element of  $P$ , so that the projective space  $P$  parametrizes triples as above up to birational isomorphism.

Let  $W \subset \Gamma(X, \mathcal{O}(-K_X))$  be the subspace so that  $P$  is the associated projective space and  $n = \mathrm{rank} W$ . We have a natural homomorphism

$$\phi : V \otimes \mathcal{O}_{X^n} \rightarrow \bigoplus_{i=1}^n p_i^* \mathcal{O}(-K_X)$$

The divisor  $D$  where the determinant  $\det(\phi)$  vanishes parametrizes, birationally, tuples of the form  $(E, S; R; p_1, \dots, p_n)$  where the  $p_i$  are  $n$  additional points which are not subject to any additional relation. Moreover,  $D$  has trivial canonical bundle.

If  $n$  is at most the dimension of the linear system, then the parameter space is rational. But new things happen when  $n$  is larger than that, when one gets a divisor on a product of rational surfaces, in fact given by the vanishing of  $\det(\phi)$ . This moduli space  $V_n$ , say, fibers over the rational variety  $V_{n-1}$ , with the general fibre being an elliptic curve. The involution  $x \mapsto -x$  on the general fibre gives rise to an involution  $\tau$  on  $V_n$ . So  $V_n/\tau$  fibers over a rational variety with  $\mathbb{P}^1$  fibres. Hence  $V_n/\tau$  is unirational.

## 2. THE MODULAR VARIETIES OF INTEREST

In the context of the congruence subgroup  $\Gamma_1(N)$  of  $\mathrm{SL}(2, \mathbb{Z})$ , the elliptic modular varieties  $V$  with non-positive canonical bundles are associated with pairs  $(N, k)$  such that there is at most one modular form of level  $N$  and weight  $k$ . The complete list of such pairs is given in the following table:

$N$	$k$
1	$\leq 23, 25, 26$ , odd
2	$\leq 11$ , odd
3	$\leq 8$
4	$\leq 6$
5, 6	$\leq 4$
= 7, 8	$\leq 3$
9, 10, 11, 12, 14, 15	2

One can similarly make the corresponding tables for the groups  $\Gamma_0(N)$  and  $\Gamma_0(N^2) \cap \Gamma_1(N)$ .

## 3. THE CALABI-YAU 11-FOLD ASSOCIATED TO $\Delta$

$\Delta$  is a generator of  $S_{12}(\mathrm{SL}(2, \mathbb{Z}))$ . The object is to show that the Kuga-Sato variety  $\mathcal{E}^{(10)}$  is birational to an eleven-dimensional Calabi-Yau variety  $V$ . We will use  $\equiv$  to denote birational equivalence. It is easy to see that for any  $r \geq 0$ ,

$$\mathcal{E}^r \equiv \mathcal{M}_1(r+1),$$

where  $\mathcal{M}_g(k)$  is the moduli space genus  $g$  curves with  $k$  marked points, with compactification  $\overline{\mathcal{M}}_g(k)$ .



So we need to find a *birational model*  $V$  of  $\overline{\mathcal{M}}_1(11)$  such that  $V$  is Calabi-Yau. Let  $S$  be the surface obtained by *blowing up* 4 *general points*  $P_1, P_2, P_3, P_4$  in  $\mathbb{P}^2$ . Let  $E$  be an elliptic curve with  $n+5$  general points  $Q_0, Q_1, \dots, Q_{n+4}$ , and use  $|3Q_0|$  to define a morphism  $E \rightarrow \mathbb{P}^2$ .

Using an automorphism of  $\mathbb{P}^2$  we may assume:  $Q_i = P_i$  for  $1 \leq i \leq 4$ . The embedding  $E \rightarrow \mathbb{P}^2$  lifts to a morphism  $\varphi : E \rightarrow S$ , and the *adjunction formula* gives  $\varphi(E) \in |\mathcal{K}_S^{-1}|$ .

Get a *rational map*  $\mathcal{M}_1(n+5) \rightarrow S^n$ ,

$$(E, \{Q_0, \dots, Q_{n+4}\}) \rightarrow (P_5, \dots, P_{n+4}).$$

$W := \Gamma(S, \mathcal{K}_S^{-1})$  has dim. 6, and  $\exists$  a hom (of sheaves on  $S^n$ ):

$$f_n : W \otimes \mathcal{O}_{S^n} \rightarrow \mathcal{K}_S^{-1} \boxtimes \dots \boxtimes \mathcal{K}_S^{-1}.$$

$\text{Ker}(f_n)$  is the vector space of *sections in  $W$  vanishing at*  $= P_5, \dots, P_{n+4}$ . The associated projective space then identifies with the collection of all (general) points in  $\mathcal{M}_1(11)$  giving rise to this point on  $S^n$ .

Put

$$V_n := \text{Proj}_{S^n}(\text{coker}({}^t f_n))$$

where

$${}^t f_n : \mathcal{K}_S \boxtimes \dots \boxtimes \mathcal{K}_S \rightarrow W^\vee \otimes \mathcal{O}_S^\vee.$$

Then  $V_n$  is birational to  $\mathcal{M}_1(n+5)$ . We have  $\text{corank}({}^t f_n) = 6 - n$  at a general point of  $S^n$ . And there is a natural map

$$\pi : V_n \rightarrow S^n$$

**n ≤ 5:**  $\pi$  is surjective with fibres  $\mathbb{P}^{5-n}$ . Hence  $V_n$ , which is  $\equiv \overline{\mathcal{M}}_1(n+5)$ , is a *rational variety* in this case. Note that  $V_5$  is just  $S^5$ .

**n = 6:**  $V = V_6$  is a (reduced) *divisor in  $S^6$* , hence Gorenstein. It is defined by the vanishing of  $\det(f_n)$ , which is a section of  $\mathcal{K}_{S^n}^{-1}$ . So  $\mathcal{K}_V$  is **trivial**. We already know that  $h^{(11,0)} = 1$  and  $h^{(p,0)} = 0$  for  $0 < p < 11$  for  $\mathcal{E}^{10}$ . These also hold for  $V$ . So  $V$  is Calabi-Yau. Also, the whole construction is rationally defined.

$V = V_6$  fibers over  $V_5$  with fibres of dimension 1, corresponding to cubics passing through 10 points. The natural involution on the general fibre, which is an elliptic curve, gives rise to one, call it  $\tau$ , on  $V$ . The quotient  $V/\tau$  is unirational because it is a family of rational curves on a smooth rational surface. Clearly,  $\tau$  must act by  $-1$  on the one dimensional space  $H^0(V, \Omega^{11})$ .

□

## 4. A C-Y 7-FOLD OCCURRING IN LEVEL 2

Let  $E$  be an elliptic curve with origin  $o \in E$ ,  $x \in E$  a point of order 2 and  $y, z \in E$  some other (general) points. Under the morphism  $a : E \rightarrow |2[o] + [y]|$ , the divisors  $2[o] + [y]$  and  $2[x] + [y]$  are linear sections. Of the four points  $o, x, y$  and  $z$  we may assume (under the hypothesis of generality on  $y$  and  $z$ ) that no three are collinear. Thus, we can identify these with the points  $(0 : 0 : 1)$ ,  $(1 : 0 : 0)$ ,  $(0 : 1 : 0)$  and  $(1 : 1 : 1)$  respectively in order to identify  $|2[o] + [y]|$  with  $\mathbb{P}^2$ .

Conversely, let  $E$  be a cubic curve in  $\mathbb{P}^2$  which has the following properties:

- (1)  $E$  passes through the points  $(0 : 0 : 1)$ ,  $(0 : 1 : 0)$ ,  $(1 : 0 : 0)$  and  $(1 : 1 : 1)$ .
- (2) The line  $Y = 0$  is tangent to  $E$  at the point  $(0 : 0 : 1)$ .
- (3) The line  $Z = 0$  is tangent to  $E$  at the point  $(0 : 1 : 0)$ .

Then  $E$  is a curve of genus 1 for which we take  $o = (0 : 0 : 1)$  as the origin of a group law. Let  $x = (0 : 1 : 0)$ ,  $y = (1 : 0 : 0)$  and  $z = (1 : 1 : 1)$ . Then we obtain the relation

$$2[o] + [y] \simeq 2[x] + [y]$$

It follows that  $2x = o$ . Thus we have obtained  $(E, o, x, y, z)$  of the type we started with.

Direct calculation shows that the linear system of cubics in  $\mathbb{P}^2$  that satisfy the conditions above is the linear span of  $X^2Y - XYZ$ ,  $X^2Z - XYZ$ ,  $Y^2Z - XYZ$  and  $YZ^2 - XYZ$ .

Here is an **alternate construction in level 2**:

Let  $E$  be an elliptic curve with origin  $o \in E$ ,  $x \in E$  a point of order 2 and  $y, z \in E$  some other points. The morphism  $a : E \rightarrow |2[o]|$  has fibres  $2[o]$ ,  $[y] + [-y]$  and  $[z] + [-z]$  which we map to 0, 1 and  $\infty$  respectively in order to identify  $|2[o]|$  with  $\mathbb{P}^1$ . The morphism  $b : E \rightarrow |[o] + [x]|$  has fibres  $[o] + [x]$ ,  $[y] + [x - y]$  and  $[z] + [x - z]$  which we map to 0, 1 and  $\infty$  in order to identify  $|[o] + [x]|$  with  $\mathbb{P}^1$ . Thus we obtain a morphism  $a \times b : E \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$  which is constructed canonically from the data  $(E, o, x, y, z)$ .

Conversely let  $E$  be a curve of type  $(2, 2)$  in  $\mathbb{P}^1 \times \mathbb{P}^1$  which has the following properties:

- (1)  $E$  passes through the points  $(0, 0)$ ,  $(1, 1)$  and  $(\infty, \text{infy})$ .
- (2) The line  $\{0\} \times \mathbb{P}^1$  is tangent to  $E$  at the point  $(0, 0)$ .
- (3) If  $(u, 0)$  is the residual point of intersection of  $E$  with  $\mathbb{P}^1 \times \{0\}$ , then  $\{u\} \times \mathbb{P}^1$  is tangent to  $E$  at this point.

Then  $E$  is a curve of genus 0 for which we take  $o = (0, 0)$  as the origin in a group law. Let  $x = (u, 0)$ ,  $y = (1, 1)$  and  $z = (\infty, \text{infy})$ . We obtain the identities  $2[o] \simeq 2[x]$  from which it follows that  $2x = o$ . Thus we have recovered the data  $(E, o, x, y, z)$ .

### 5. REMARK ON ELLIPTIC CURVES WITH LEVEL 3 STRUCTURE

Let  $E$  be an elliptic curve with origin  $o \in E$ ,  $x \in E$  a point of order 3 and  $y \in E$  some other point. The morphism  $a : E \rightarrow |2[o]|$  has fibres  $2[o]$ ,  $[x] + [2x]$  and  $[y] + [-y]$  which we map to 0, 1 and  $\infty$  respectively in order to identify  $|2[o]|$  with  $\mathbb{P}^1$ . The morphism  $b : E \rightarrow |[o] + [x]|$  has fibres  $[o] + [x]$ ,  $2[2x]$  and  $[y] + [x - y]$  which we map to 0, 1 and  $\infty$  respectively in order to identify  $|[o] + [x]|$  with  $\mathbb{P}^1$ . Thus we obtain a morphism  $a \times b : E \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$  which is constructed canonically from the data  $(E, o, x, y)$ .

Conversely, let  $E$  be a curve of type  $(2, 2)$  in  $\mathbb{P}^1 \times \mathbb{P}^1$  which has the following properties:

- (1)  $E$  is tangent to the line  $\{0\} \times \mathbb{P}^1$  at the point  $(0, 0)$ .
- (2)  $E$  is tangent to the line  $\mathbb{P}^1 \times \{1\}$  at the point  $(1, 1)$ .
- (3)  $E$  passes through the points  $(1, 0)$  and  $(\infty, \infty)$ .

Then  $E$  is a curve of genus 1 and we use  $o = (0, 0)$  as the origin of a group law on  $E$ . Let  $x = (1, 0)$ ,  $y = (\infty, \infty)$  and  $p = (1, 1)$ . We obtain the identities,

$$2[o] \simeq [x] + [p] \qquad [o] + [x] \simeq 2[p]$$

It follows that  $p = 2x$  and  $3x = o$ . We have thus recovered the data  $(E, o, x, y)$  that we started with.

Direct calculations shows that the linear system of cubic curves in  $\mathbb{P}^2$  satisfying the conditions given above is the linear span of  $X^2Y - XYZ$ ,  $Y^2Z - XYZ$  and  $Z^2X - XYZ$ .

### 6. LEVEL 3 AND A CY 5-FOLD

We will construct a 5-fold with trivial canonical bundle and singularities only in dimension 2 or less such that its middle cohomology represents the motive of the (unique) modular form of level 3 and weight 6.

Consider the linear system  $P$  of cubics in  $\mathbb{P}^2$  that is spanned by the curves  $X^2Y - XYZ$ ,  $Y^2Z - XYZ$  and  $Z^2X - XYZ$ ; this system is stable under the cyclic automorphism  $X \rightarrow Y \rightarrow Z \rightarrow X$  of  $\mathbb{P}^2$ . Each curve in the linear system  $P$  is tangent to the line  $Z = 0$  at the point

$p_Y = (0 : 1 : 0)$ ; similarly, the curve is tangent to  $X = 0$  at the point  $p_Z = (0 : 0 : 1)$  and to  $Y = 0$  at the point  $p_X = (1 : 0 : 0)$ . Moreover, each curve passes through the point  $p_0 = (1 : 1 : 1)$ . We note that the linear system  $P$  is precisely the collection of cubic curves in  $\mathbb{P}^2$  that satisfy these conditions.

In the divisor class group of a smooth curve in this linear system we obtain the identities

$$2p_Y + p_X = 2p_Z + p_Y = 2p_X + p_Z = p_X + p_0 + r$$

where  $r$  denotes the remaining point of intersection of the curve with the line  $Y = Z$  that joins  $p_X$  and  $p_0$ . In particular, we note that  $p_Y - p_X$  is of order 3 in this class group and  $p_Z - p_X = 2(p_Y - p_X)$ .

Conversely, suppose we are given a smooth curve  $E$  of genus 1 and a line bundle  $\xi$  of order 3 on  $E$ ; moreover, suppose that three distinct points  $p, q$  and  $r$  are marked on  $E$ . We then obtain two additional points  $a$  and  $b$  on  $E$  such that  $a - p = \xi$  and  $b - p = 2\xi$  in the divisor class group of  $E$ . Consider the morphism  $E \rightarrow \mathbb{P}^2$  that is given by the linear system of the divisor  $p + q + r$ . Moreover, we choose co-ordinates on  $\mathbb{P}^2$  so that the point  $p$  goes to  $p_X, q$  goes to  $p_0, a$  goes to  $p_Y$  and  $b$  goes to  $p_Z$ . This gives an embedding of  $E$  as a curve in  $\mathbb{P}^2$  that belongs to the linear system  $P$ .

Let  $S$  denote the surface obtained by blowing up  $\mathbb{P}^2$  at the four points  $p_X, p_Y, p_Z$  and  $p_0$ , and then further blowing up the resulting surface at the ‘‘infinitely near points’’ that correspond to  $Z = 0$  at  $p_Y$ , to  $X = 0$  at  $p_Z$  and to  $Y = 0$  at  $p_X$ . Let  $H$  denote the inverse image in  $S$  of a general line in  $\mathbb{P}^2$ ; let  $E_X, E_Y, E_Z$  and  $E_0$  denote the strict transforms of the exceptional loci of the first blow-up over the points  $p_X, p_Y, p_Z$  and  $p_0$  respectively; let  $F_X, F_Y$  and  $F_Z$  denote the exceptional divisors of the second blow-up. The anti-canonical divisor  $-K_S = 3H - E_X - E_Y - E_Z - E_0 - 2(F_X - F_Y - F_Z)$  has a base-point free complete linear system  $|-K_S|$  which can be identified with  $P$ . Let  $T$  denote the natural incidence locus in  $S \times P$ . The variety

$$X = T \times_P T \times_P T$$

is a singular 5-fold which is Gorenstein and has trivial canonical bundle. Moreover, an open subset of  $X_0$  parametrizes tuples of the form  $(E, \xi, p, q, r, s, t, u)$  where  $E$  is a curve of genus 1,  $\xi$  is a line bundle of order 3 on  $E$  and  $p, q, r, s, t$  and  $u$  are six distinct points on  $E$ .

Let  $L_X, L_Y, L_Z$  denote the strict transforms in  $S$  of the lines in  $\mathbb{P}^2$  defined by  $X = 0, Y = 0, Z = 0$  respectively. Further, let  $R$  be the strict transform in  $S$  of the curve in  $\mathbb{P}^2$  defined by

$$X^2Z + Y^2X + Z^2Y - 3XYZ = 0$$

This is the unique cubic in  $\mathbb{P}^2$  that has a node at  $p_0$  and is tangent to  $X = 0$  at  $p_Y$ , to  $Y = 0$  at  $p_Z$  and to  $Z = 0$  at  $p_X$ . It follows that  $R$  is a smooth rational curve that meets  $E_0$  in a pair of distinct points and the triple  $(R, L_X, E_Y)$  (respectively  $(R, L_Y, E_Z)$  and  $(R, L_Z, E_X)$ ) consists of smooth curves that meet pairwise transversally.

The morphism  $S \rightarrow P^*$  induced by the linear system  $P$  can be factorized via a double cover  $S \rightarrow W$  which is ramified along  $R$ . Each of the curves  $L_X$  and  $E_Y$  (respectively  $L_Y$  and  $E_Z$ ;  $L_Z$  and  $E_X$ ) is mapped isomorphically onto the same smooth irreducible curve  $G_Z$  (respectively  $G_X$ ;  $G_Y$ ) in  $W$ ; the curve  $R$  is mapped isomorphically onto the branch locus  $Q$  in  $W$ . The morphism  $W \rightarrow P^*$  collapses the curves  $G_X$  (respectively  $G_Y$  and  $G_Z$ ) to a point  $q_X$  (respectively  $q_Y$  and  $q_Z$ ) in  $P^*$ ; in fact  $W \rightarrow P^*$  is identified with the blow-up of  $P^*$  at these points. Moreover,  $Q$  is mapped to a plane quartic  $\bar{Q}$  which has cusps at these three points.

Let  $T$  denote the incidence locus in  $P \times S$  as above. It is the pull-back via  $S \rightarrow P^*$  of the natural incidence locus  $I \subset P \times P^*$ . The latter can be identified (via the projection  $I \rightarrow P^*$ ) with the projective bundle of 1-dimensional linear subspaces of the tangent bundle of  $P^*$ . Hence, the exceptional curve  $G_X$  (respectively  $G_Y$  and  $G_Z$ ) of the blow-up  $W \rightarrow P^*$  can be identified with the fibre  $I_X$  of  $I \rightarrow P^*$  over the point  $q_X$  (respectively  $q_Y$  and  $q_Z$ ). Thus we obtain natural maps  $E_\alpha \rightarrow T$  and  $L_\alpha \rightarrow T$  that are sections of the  $\mathbb{P}^1$ -bundle  $T \rightarrow S$  over the curves  $E_\alpha$  and  $L_\alpha$  respectively; let  $\tilde{E}_\alpha$  and  $\tilde{L}_\alpha$  denote the images. Let  $T_X$  (respectively  $T_Y$ ;  $T_Z$ ) denote the fibre of  $T$  over the point of intersection of  $L_Y$  and  $E_Z$  (respectively  $L_Z$  and  $E_X$ ;  $L_X$  and  $E_Y$ ).

The tangent direction along  $\bar{Q}$  gives a rational morphism (defined outside the cusps) from  $\bar{Q}$  to  $I$ . It follows that this extends to a section  $R \rightarrow T$  of  $T \rightarrow S$  over  $R$  and gives a curve  $\tilde{R}$  in  $T$ . The quadruple of curves  $\tilde{R}, T_X, \tilde{L}_Y, \tilde{E}_Z$  (respectively,  $\tilde{R}, T_Y, \tilde{L}_Z, \tilde{E}_X$ ;  $\tilde{R}, T_Z, \tilde{L}_X, \tilde{E}_Y$ ) meet pairwise transversally in a single point  $r_X$  (respectively  $r_Y$ ;  $r_Z$ ) in  $T$ . The curve  $\tilde{R}$  in  $T$  is mapped to a nodal cubic  $\bar{R}$  in  $P$  for which  $I_X, I_Y$  and  $I_Z$  are inflectional tangents. The curves  $T_X, \tilde{L}_Y, \tilde{E}_Z$  (respectively  $T_Y, \tilde{L}_Z, \tilde{E}_X$ ;  $T_Z, \tilde{L}_X, \tilde{E}_Y$ ) in  $T$  lie over  $I_X$  (respectively  $I_Y$ ;  $I_Z$ ) in  $P$ .

The singular locus of the morphism  $T \rightarrow P$  consists of the curves  $\tilde{R}, T_\alpha, \tilde{L}_\alpha$  and  $\tilde{E}_\alpha$  for  $\alpha = X, Y, Z$  as described above.

The singular fibres of  $T \rightarrow P$  then have the following description:

- (1) If  $a$  is a smooth point of  $\bar{R}$  which is not a point of inflection then the fibre  $C_a$  is a rational curve in  $S$  with a single ordinary node.

- (2) If  $b$  which is on an inflectional tangent (i. e. one of the lines  $I_X, I_Y, I_Z$ ) of  $\overline{R}$  but is *not* a point of inflection of  $\overline{R}$  then the fibre  $C_b$  is a curve with three components and three nodes (i. e. a “triangle” of  $\mathbb{P}^1$ 's).
- (3) If  $c$  is a point of inflection of the curve  $\overline{R}$ , then  $C_c$  consists of three  $\mathbb{P}^1$ 's that pass through a point and (since  $C_c$  lies on a smooth surface  $S$ ) is locally a complete intersection.
- (4) If  $d$  is the node of  $\overline{R}$  then the fibre  $C_d$  consists of a pair of smooth  $\mathbb{P}^1$ 's in  $S$  that meet in a pair of points. In fact the curves are  $E_0$  and the strict transform in  $S$  of the curve in  $\mathbb{P}^2$  defined by the equation

$$X^2Y + Y^2Z + Z^2X - 3XYZ = 0$$

In particular, the elliptic fibration  $T \rightarrow P$  is semi-stable but for the three fibres over the points of inflection of  $\overline{R}$ .

Now consider the variety  $X = T \times_P T \times_P T$ . The singular points of  $X_0$  consist of triples  $(x, y, z)$  of points of  $T$ , where at least two of these points are critical points for the morphism  $T \rightarrow P$ . In particular, these points lie over the union of  $R$  and  $I_X, I_Y$  and  $I_Z$ . Since the singular points of each of the fibres described above are isolated, it follows that the singular locus of  $X$  has components of dimension at most 2.

## 7. LEVEL 4 AND A C-Y 5-FOLD

Let  $E$  be an elliptic curve with origin  $o \in E$ ,  $x \in E$  a point of order 4. Under the morphism  $a : E \rightarrow |\mathbb{P}^1|$ , the divisors  $3[o]$ ,  $2[2x] + [o]$ ,  $[2x] + 2[x]$  and  $[o] + [x] + [3x]$  are linear sections of the image curve. Let  $p$  denote the point of intersection of the lines corresponding to  $3[o]$  and  $[2x] + 2[x]$ . No three of the points  $o, 2x, 3x$  and  $p$  are collinear. Thus we can identify  $|\mathbb{P}^1|$  with  $\mathbb{P}^2$  in such a way that  $o$  is identified with  $(0 : 1 : 0)$ ,  $3x$  is identified with  $(0 : 0 : 1)$ ,  $p$  is identified with  $(1 : 0 : 0)$  and  $2x$  is identified with  $(1 : 1 : 1)$ . Thus we obtain a morphism  $a : E \rightarrow \mathbb{P}^2$  which is constructed canonically from the data  $(E, o, x)$ .

Conversely, let  $E$  be a cubic curve in  $\mathbb{P}^2$  which has the following properties:

- (1)  $E$  passes through the points  $(0 : 1 : 0)$ ,  $(0 : 0 : 1)$ ,  $(1 : 1 : 1)$  and  $(1 : 0 : 1)$ .
- (2) The line  $Z = 0$  is an inflectional tangent to  $E$  (at the point  $(0 : 1 : 0)$ ).

- (3) The line  $X = 0$  is tangential to the curve  $E$  at the point  $(0 : 0 : 1)$ .
- (4) the line  $Y = 0$  is tangential to the curve  $E$  at the point  $(1 : 0 : 1)$ .

Then  $E$  is a curve of genus 0 and we use  $o = (0 : 1 : 0)$  as the origin of a group law on  $E$ . Let  $x = (1 : 0 : 1)$ ,  $p = (0 : 0 : 1)$  and  $q = (1 : 1 : 1)$ . We obtain the identities

$$3[o] \simeq [o] + 2[p] \qquad \simeq [p] + 2[x] \simeq [o] + [q] + [x]$$

It follows that  $2p = o$ ,  $2x = p$  and  $q = -x = 3x$ . Thus we have recovered the data  $(E, o, x)$  that we started with.

Direct calculation shows us that the linear system of cubic curves in  $\mathbb{P}^2$  that satisfy the above conditions is the linear span of  $YZ(Y - Z)$  and  $X(X - Z)^2$ .

Here is an **alternate construction in level 4**:

Let  $E$  be an elliptic curve with origin  $o \in E$ ,  $x \in E$  a point of order 4 and  $y \in E$  some other point. The morphism  $a : E \rightarrow |2[o]|$  has fibres  $2[o]$ ,  $2[2x]$  and  $[y] + [-y]$  which we map to 0, 1 and  $\infty$  respectively in order to identify  $|2[o]|$  with  $\mathbb{P}^1$ . The morphism  $b : E \rightarrow |[o] + [x]|$  has fibres  $[o] + [x]$ ,  $[2x] + [3x]$  and  $[y] + [x - y]$  which we map to 0, 1 and  $\infty$  respectively in order to identify  $|[o] + [x]|$  with  $\mathbb{P}^1$ . Thus we obtain a morphism  $a \times b : E \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$  which is constructed canonically from the data  $(E, o, x, y)$ .

Conversely, let  $E$  be a curve of type  $(2, 2)$  in  $\mathbb{P}^1 \times \mathbb{P}^1$  which has the following properties:

- (1)  $E$  is tangent to the line  $\{0\} \times \mathbb{P}^1$  at the point  $(0, 0)$ .
- (2)  $E$  is tangent to the line  $\{1\} \times \mathbb{P}^1$  at the point  $(1, 1)$ .
- (3) If  $E$  meets  $\mathbb{P}^1 \times \{0\}$  at  $(0, 0)$  and  $(u, 0)$  and  $E$  meets  $\mathbb{P}^1 \times \{1\}$  at  $(1, 1)$  and  $(v, 1)$ ; then  $u = v$ .
- (4)  $E$  passes through the point  $(\infty, \infty)$ .

Then  $E$  is a curve of genus 0 and we use  $o = (0, 0)$  as the origin of a group law on  $E$ . Let  $x = (u, 0)$ ,  $p = (v, 1)$  and  $q = (1, 1)$ . We obtain the identities,

$$\begin{aligned} 2[o] &\simeq 2[q] & [o] + [x] &\simeq [p] + [q] \\ 2[0] &\simeq [x] + [p] & & \text{(from condition 3 above)} \end{aligned}$$

It follows that  $q = 2x$ ,  $p = 3x$  and  $4x = o$ . Let  $y = (\infty, \text{infy})$ . We have thus recovered the data  $(E, o, x, y)$  that we started with.

## 8. FORMS OF WEIGHT 4 AND CALABI-YAU THREEFOLDS

Let  $E$  be an elliptic curve with  $o \in E$  as its origin and  $x \in E$  a point of order 5. Under the morphism  $a : E \rightarrow |3[o]|$ , the divisors  $3[o]$ ,  $[o] + [x] + [4x]$ ,  $2[x] + [3x]$  and  $2[3x] + [4x]$  are linear sections. There is a unique identification of  $|3[o]|$  with  $\mathbb{P}^2$  under which these sections are identified with  $Z = 0$ ,  $X = 0$ ,  $X + Y + Z = 0$  and  $Y = 0$  respectively.

Conversely, let  $E$  be a cubic curve in  $\mathbb{P}^2$  which has the following properties:

- (1)  $E$  passes through the points  $(0 : 1 : 0)$ ,  $(0 : 1 : -1)$ ,  $(1 : 0 : -1)$  and  $(0 : 0 : 1)$ .
- (2) The line  $Z = 0$  is an inflectional tangent to  $E$  (at the point  $(0 : 1 : 0)$ ).
- (3) The line  $X + Y + Z = 0$  is tangent to  $E$  at the point  $(1 : 0 : -1)$ .
- (4) The line  $Y = 0$  is tangent to  $E$  at the point  $(0 : 1 : -1)$ .

The  $E$  is a curve of genus 1. Let  $o = (0 : 1 : 0)$ ,  $x = (0 : 1 : -1)$ ,  $p = (0 : 0 : 1)$  and  $q = (1 : 0 : -1)$ . We use  $o$  as the origin of the group law on  $E$ . We obtain the identities,

$$3[o] \simeq [o] + [x] + [p] \simeq 2[x] + [q] \simeq 2[q] + [p].$$

It follows that  $q = 2x$ ,  $p = 4x$  and  $5x = o$ . Thus we have obtained the data  $(E, o, x)$  that we started with.

Direct calculation shows us that the linear system of cubics in  $\mathbb{P}^2$  satisfying the above conditions is the linear span of  $YZ(X + Y + Z)$  and  $YZ(Y + Z) - X(X + Z)^2$ .

Let  $E$  be an elliptic curve with  $o \in E$  as its origin and  $x \in E$  a point of order 5. The morphism  $a : E \rightarrow |2[o]|$  has fibres  $2[o]$ ,  $[x] + [4x]$  and  $[2x] + [3x]$ , which we map to 0, 1 and  $\infty$  to identify  $|2[o]|$  with  $\mathbb{P}^1$ . Similarly, the morphism  $b : E \rightarrow |2[x]|$  has fibres  $2[x]$ ,  $[o] + [2x]$  and  $[3x] + [4x]$ , which we map to 0, 1 and  $\infty$  to identify  $|2[x]|$  with  $\mathbb{P}^1$ . Thus we obtain a morphism  $a \times b : E \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ , which is constructed canonically from the data  $(E, o, x)$ .

Conversely, let  $E$  be a curve of type  $(2, 2)$  in  $\mathbb{P}^1 \times \mathbb{P}^1$  which is

- (1) tangent to  $\mathbb{P}^1 \times \{0\}$  at the point  $(1, 0)$ ,
- (2) tangent to  $\{0\} \times \mathbb{P}^1$  at the point  $(0, 1)$ ,
- (3) passes through the points  $(\infty, 1)$ ,  $(\infty, \infty)$ , and  $(1, \infty)$ .

Then  $E$  is a curve of genus 1. Let  $o = (0, 1)$  and  $x = (1, 0)$ , which are points on  $E$ . We use  $o$  as the origin of the group law on  $E$ . Let  $p$ ,  $q$ ,  $r$  denote the points  $(\infty, 1)$ ,  $(\infty, \infty)$  and  $(1, \infty)$  respectively. We obtain



the identities,

$$\begin{aligned} 2[o] &\simeq [p] + [q] & 2[x] &\simeq [q] + [r] \\ 2[o] &\simeq [x] + [r] & 2[x] &\simeq [o] + [p] \end{aligned}$$

We solve these to show that  $a = 2x$ ,  $b = 3x$ ,  $c = 4x$ , and  $5x = o$ . We have thus recovered the data  $(E, o, x)$  that we started with.

Now, appealing to the fibre product paper of Schoen ([?]), we can deduce that our canonical object is Calabi-Yau.

### 9. CM FORMS AND C-Y VARIETIES OVER SUITABLE EXTENSIONS

Let  $E$  be an elliptic curve and  $n \geq 1$ . Put

$$B := \{x \in E^{n+1} \mid \sum_{j=1}^{n+1} x_j = 0\},$$

which admits an action by the alternating group  $A_{n+1}$ . Consider the quotient

$$X := B/A_{n+1}.$$

The following result is proved in [PR], where the smoothness of the model for  $n = 3$  appeals to ideas of Cynk and Hulek.

**Theorem**  *$X$  has trivial canonical bundle, with  $H^0(X, \Omega_X^p) = 0$  if  $0 < p < n$ . If  $n \leq 3$ , there is a smooth model  $\tilde{X}$  which is Calabi-Yau.*

A submotive  $M$  of rank 4 splits off of  $H^3(\tilde{X})$  corresponding to  $\text{sym}^3(H^1(E))$ , of Hodge type  $\{(3, 0), (2, 1), (1, 2), (0, 3)\}$ . It is simple iff  $E$  is not of CM type, and in this case  $\tilde{X}$  is not rigid.

Here is a sketch of proof of Theorem 2. Let  $\Psi$  be the Hecke character of an imaginary quadratic field  $K$  attached to  $f$ , so that  $L(s, f) = L(s, \Psi)$ . Pick an algebraic Hecke character  $\lambda$  of  $K$  of weight 1. Then  $\Psi/\lambda^{k-1}$  is a finite order character  $\nu$ . Attach a Calabi-Yau to  $\lambda^{k-1}$  by using the theorem above. Vary  $\lambda$ .

□

We will discuss Theorem 3 elsewhere in detail, where for the key case  $m = 3$  is partly understood via the descent to  $\text{GSp}(4)/\mathbb{Q}$  ([RS]).

## 10. BEHAVIOR UNDER TAKING PRODUCTS

Suppose  $X_i$  is a double cover of smooth variety  $Y_i$  branched along a smooth divisor  $D_i$  for  $i = 1, 2$ ; let  $\iota_i$  denote the associated involutions on  $X_i$ .

The product variety  $X_1 \times X_2$  carries an action of the involution  $\iota = \iota_1 \times \iota_2$  which has  $D_1 \times D_2$  as its fixed locus. This is a codimension two transverse intersection of the divisors  $D_1 \times X_2$  and  $X_1 \times D_2$ .

Let  $X_{12}$  be the blow-up of  $X_1 \times X_2$  along  $D_1 \times D_2$  and  $E_{12}$  be the exceptional divisor. Then  $E_{12}$  is isomorphic to the projective bundle over  $D_1 \times D_2$  associated with the rank two vector bundle  $L_1 \oplus L_2$ , where  $L_i = p_i^* N_{D_i/X_i}$ . The strict transform  $E_1$  of  $D_1 \times X_2$  (respectively  $E_2$  of  $X_1 \times D_2$ ) in  $X_{12}$  is a divisor that meets  $E_{12}$  in a section of this projective bundle; the two intersections  $E_1 \cap E_{12}$  and  $E_2 \cap E_{12}$  are disjoint.

Since  $D_1 \times D_2$  is the fixed locus of  $\iota$ , this involution lifts to  $X_{12}$ ; we denote this lift also by  $\iota$  by abuse of notation. The (scheme-theoretic) fixed locus of  $\iota$  on  $X_{12}$  is the smooth divisor  $E_{12}$  and hence the quotient  $Z_{12}$  of  $X_{12}$  by this involution is a smooth variety. In other words,  $X_{12}$  is a double cover of  $Z_{12}$  branched along the smooth divisor  $D_{12}$  which is the image of  $E_{12}$ .

The involution  $\iota_1 \times id_{X_2}$  also lifts to  $X_{12}$  since the base of the blow-up is contained in its fixed locus. Moreover, it commutes with  $\iota$  and hence descends to an involution  $\tau$  on  $Z_{12}$ . The (scheme-theoretic) fixed locus of this involution on  $Z_{12}$  is the *disjoint* union of the images of  $E_1$  and  $E_2$  and is thus again a smooth divisor. Thus  $Z_{12}$  is the double cover of a variety  $Y_{12}$  branched along a smooth divisor. Moreover, one checks that  $Y_{12}$  is just the blow-up of  $Y_1 \times Y_2$  along  $D_1 \times D_2$  (where by abuse of notation we are using the same notation for the divisors  $D_i$  in  $X_i$  and for their images in  $Y_i$ ).

Finally, let us calculate the canonical bundle of  $Z_{12}$  varieties. We note that the double cover  $X$  of smooth variety  $Y$  branched along a smooth divisor  $D$  is obtained by choosing an isomorphism of  $\mathcal{O}_Y(D)$  with the square of some line bundle  $L$  on  $Y$ . In this case, the canonical bundle of  $X$  is the pull back of  $K_Y \otimes L$ .

Let  $L_i$  denote the square root of  $\mathcal{O}_{Y_i}(D_i)$ . The canonical bundle of  $K_{X_i}$  is the pull-back of  $K_{Y_i} \otimes L_i$ ; so if  $X_i$  is a Calabi-Yau variety, then  $K_{Y_i}$  is the dual of  $L_i$ . If  $E$  denotes the exceptional locus of the morphism  $e : Y_{12} \rightarrow Y_1 \times Y_2$ , then we see that if  $\hat{D}_1$  denotes the strict transform of  $D_1 \times Y_2$  in  $Y_{12}$  we have

$$\mathcal{O}_{Y_{12}}(\hat{D}_1 + E) = e^* \mathcal{O}_{Y_1 \times Y_2}(D_1 \times Y_2)$$

Similarly, we have

$$\mathcal{O}_{Y_{12}}(\hat{D}_2 + E) = e^* \mathcal{O}_{Y_1 \times Y_2}(Y_1 \times D_2)$$

It follows that

$$\mathcal{O}_{Y_{12}}(\hat{D}_1 + \hat{D}_2) = e^*(L_1 \otimes L_2 \otimes \mathcal{O}_{Y_{12}}(-E))^2$$

Since the canonical bundle of  $Y_{12}$  is

$$K_{Y_{12}} = e^* K_{Y_1 \times Y_2} \otimes \mathcal{O}_{Y_{12}}(E) = e^*(L_1 \boxtimes L_2)^{-1} \otimes \mathcal{O}_{Y_{12}}(E)$$

it follows that the canonical bundle of the double cover  $Z_{12}$  is trivial.

Theorem 4 follows from applying this construction. □

APPENDIX: A CONSEQUENCE OF THE HIRZEBRUCH  
RIEMANN-ROCH THEOREM

**Theorem** *The Hirzebruch Riemann-Roch theorem does not impose any restrictions on the Euler characteristic of an odd-dimensional smooth and projective variety of dimension at least 3.*

**Proof** Let  $X$  be any smooth projective variety. Recall that the Todd classes of a variety are the multiplicative classes defined by the generating function

$$\text{td}(t) = \frac{t}{1 - \exp(-t)} = 1 + \frac{t}{2} + \frac{t^2}{12} - \frac{t^4}{720} + \frac{t^6}{30240} + O(t^8)$$

Fix an integer  $m$  and for  $i = 1, \dots, m$ , let  $\beta_i$  be algebraic numbers such that

$$\text{td}(t) \equiv \prod_{i=1}^m (1 + \beta_i t) \pmod{t^{m+1}}$$

For  $X$  of dimension  $m$ , let  $c_i$  for  $i = 1, \dots, m$  be the Chern classes of its tangent bundle. Let  $\gamma_i$  for  $i = 1, \dots, m$  be the Chern roots so that

$$1 + c_1 t + c_2 t^2 + \dots + c_m t^m = \prod_{i=1}^m (1 + \gamma_i t)$$

The Todd polynomial of the variety is then given by

$$\text{Todd}(t) = \prod_{i=1}^m \text{td}(\gamma_i t)$$

The coefficient of  $t^m$  in  $\text{Todd}(t)$  is the  $m$ -th Todd class  $\text{Todd}_m$  of the variety. Making use of the above expression  $\text{td}(t) \pmod{t^{m+1}}$ ,

$$\text{Todd}(t) \equiv \prod_{i,j=1}^m (1 + \beta_j \gamma_i t) \equiv \prod_{j=1}^m (1 + c_1(\beta_j t) + c_2(\beta_j t)^2 + \cdots + c_m(\beta_j t)^m \pmod{t^{m+1}})$$

Hence, the coefficient of  $c_m$  in the  $m$ -th Todd class  $\text{Todd}_m$  is  $\sum_{j=1}^m \beta_j$ .

We can compute this as follows. Consider the function  $f(t) = \log(\text{td}(t))$ . It has an expression modulo  $t^{m+1}$  as

$$f(t) = \sum_{j=1}^m \log(1 + \beta_j t) = \sum_{j=1}^m \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} (\beta_j t)^k \pmod{t^{m+1}}$$

On the other hand we have  $f(-t) = f(t) - t$  so that in the expression of  $f(t)$  as a power series in  $t$ , all the odd degree terms except  $t$  have coefficient 0. In particular, it follows that  $\sum_{j=1}^m \beta_j$  is 0 whenever  $m$  is odd and  $m > 1$ .

To summarize, we have proved that the coefficient of the top Chern class in the Todd class of an odd-dimensional variety is 0. Since this top Chern class can be identified with the Euler characteristic we have the result that we wished to prove.

## REFERENCES

- [Clo1] L. Clozel. Motifs et formes automorphes: applications du principe de fonctorialité. In *Automorphic forms, Shimura varieties, and L-functions, Vol. I (Ann Arbor, MI, 1988)*, volume 10 of *Perspect. Math.*, pages 77–159. Academic Press, Boston, MA, 1990.
- [Clo2] L. Clozel. Représentations galoisiennes associées aux représentations automorphes autoduales de  $\text{GL}(n)$ . *Inst. Hautes Études Sci. Publ. Math.* (1991), 97–145.
- [Del] P. Deligne. Formes modulaires et représentations  $\ell$ -adiques. *Séminaire Bourbaki 1968/69* **355** (1971), 139–172.
- [DMOS] P. Deligne, J. S. Milne, A. Ogus, and K.-y. Shih. *Hodge cycles, motives, and Shimura varieties*, volume 900 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1982.
- [ES] N. Elkies and M. Schuett. Modular forms and K3 surfaces, preprint, arxiv 0809.0830, 2008.
- [KW] C. Khare and J.-P. Wintenberger. Serre’s modularity conjecture. I. *Invent. Math.* **178** (2009), 485–504. <http://dx.doi.org/10.1007/s00222-009-0205-7>
- [Kim] H. H. Kim. Functoriality for the exterior square of  $\text{GL}_4$  and the symmetric fourth of  $\text{GL}_2$ . *J. Amer. Math. Soc.* **16** (2003), 139–183 (electronic). With Appendix 1 by Dinakar Ramakrishnan and Appendix 2 by Kim and Peter Sarnak.

- [KS] H. H. Kim and F. Shahidi. Functorial products for  $GL_2 \times GL_3$  and the symmetric cube for  $GL_2$ . *Ann. of Math. (2)* **155** (2002), 837–893. With an appendix by Colin J. Bushnell and Guy Henniart.
- [Kis] M. Kisin. Modularity of 2-adic Barsotti-Tate representations. *Invent. Math.* **178** (2009), 587–634. <http://dx.doi.org/10.1007/s00222-009-0207-5>
- [PR] K. Paranjape and D. Ramakrishnan. Quotients of  $E^n$  by  $\mathfrak{A}_{n+1}$  and Calabi-Yau manifolds. In *Algebra and number theory*, pages 90–98. Hindustan Book Agency, Delhi, 2005.
- [Ram] D. Ramakrishnan. Modularity of the Rankin-Selberg  $L$ -series, and multiplicity one for  $SL(2)$ . *Ann. of Math. (2)* **152** (2000), 45–111.
- [RS] D. Ramakrishnan and F. Shahidi. Siegel modular forms of genus 2 attached to elliptic curves. *Math. Res. Lett.* **14** (2007), 315–332.
- [YL] N. Yui and J. D. Lewis, editors. *Calabi-Yau varieties and mirror symmetry*, volume 38 of *Fields Institute Communications*, Providence, RI, 2003. American Mathematical Society.
- [YYL] N. Yui, S.-T. Yau, and J. D. Lewis, editors. *Mirror symmetry. V*, volume 38 of *AMS/IP Studies in Advanced Mathematics*, Providence, RI, 2006. American Mathematical Society.

DEPARTMENT OF MATHEMATICAL SCIENCES, IISER, MOHALI, INDIA  
*E-mail address:* kapil.iiserm@gmail.com

DEPARTMENT OF MATHEMATICSS, CALIFORNIA INSTITUTE OF TECHNOLOGY,  
PASADENA, CA 91125  
*E-mail address:* dinakar@caltech.edu