

Definable refinements of classical algebraic invariants

Martino Lupini

Università di Bologna

Caltech Logic Seminar

March 7, 2023



Funded by the European Union (ERC Stating Grant DAT, G.A. 101077154). Views and opinions expressed are however those of the author only and do not necessarily reflect those of the European Union or the European Research Council Executive Agency. Neither the European Union nor the granting authority can be held responsible for them.

Table of Contents

- 1 Motivation from topology
- 2 “Completions” of categories of algebraic-topological objects
- 3 Definable refinements of algebraic invariants
 - Finer invariants
 - Richer invariants
 - Rigid invariants

Table of Contents

- 1 Motivation from topology
- 2 “Completions” of categories of algebraic-topological objects
- 3 Definable refinements of algebraic invariants
 - Finer invariants
 - Richer invariants
 - Rigid invariants

Topology and Classification

In topology one tries to classify spaces up to **homeomorphism**

In homotopy theory the relation of **homotopy equivalence** is considered

Topology and Classification

In topology one tries to classify spaces up to **homeomorphism**

In homotopy theory the relation of **homotopy equivalence** is considered

Topology and Classification

In topology one tries to classify spaces up to **homeomorphism**

In homotopy theory the relation of **homotopy equivalence** is considered

Topology and Classification

In topology one tries to classify spaces up to **homeomorphism**

In homotopy theory the relation of **homotopy equivalence** is considered

Invariants in Algebraic Topology

One attaches to topological spaces **algebraic invariants** such as groups

(All the groups will be abelian.)

From complexes to groups

The final invariant (group) is obtained by passing via **complexes**.

Why Polish groups?

Polish: second countable, topology induced by a complete metric

Why Polish groups?

Polish: second countable, topology induced by a complete metric

The class of Polish groups:

- contains locally compact groups

Why Polish groups?

Polish: second countable, topology induced by a complete metric

The class of Polish groups:

- contains locally compact groups
- contains spaces from analysis (Banach spaces, operator algebras)

Why Polish groups?

Polish: second countable, topology induced by a complete metric

The class of Polish groups:

- contains locally compact groups
- contains spaces from analysis (Banach spaces, operator algebras)
- contains automorphism groups of “reasonable” structures

Why Polish groups?

Polish: second countable, topology induced by a complete metric

The class of Polish groups:

- contains locally compact groups
- contains spaces from analysis (Banach spaces, operator algebras)
- contains automorphism groups of “reasonable” structures
- is closed under countable products and inverse limits

Why Polish groups?

Polish: second countable, topology induced by a complete metric

The class of Polish groups:

- contains locally compact groups
- contains spaces from analysis (Banach spaces, operator algebras)
- contains automorphism groups of “reasonable” structures
- is closed under countable products and inverse limits
- is closed under closed subgroups and quotients by closed subgroups

Why Polish groups?

Polish: second countable, topology induced by a complete metric

The class of Polish groups:

- contains locally compact groups
- contains spaces from analysis (Banach spaces, operator algebras)
- contains automorphism groups of “reasonable” structures
- is closed under countable products and inverse limits
- is closed under closed subgroups and quotients by closed subgroups
- the σ -algebra of Borel sets of a Polish group is **standard** (isomorphic to the σ -algebra of Borel sets of \mathbb{R})

The homology of a Polish complex

Consider a complex of Polish groups A_* :

$$\cdots \longrightarrow A_0 \xrightarrow{\varphi_0} A_1 \xrightarrow{\varphi_1} A_2 \longrightarrow \cdots$$

The homology of a Polish complex

Consider a complex of Polish groups A_* :

$$\cdots \longrightarrow A_0 \xrightarrow{\varphi_0} A_1 \xrightarrow{\varphi_1} A_2 \longrightarrow \cdots$$

The homology of a Polish complex

Consider a complex of Polish groups A_* :

$$\dots \longrightarrow A_0 \xrightarrow{\varphi_0} A_1 \xrightarrow{\varphi_1} A_2 \longrightarrow \dots$$

In *A History of Algebraic and Differential Topology*, Dieudonné writes of

a trend that was very popular until around 1950 (although later all but abandoned), namely, to consider homology groups as topological groups for suitably chosen topologies.

The problem with cokernels

Problem: the category \mathcal{A} of abelian Polish groups is **not abelian**

Reason: a continuous group homomorphism **need not have closed image**

The problem with cokernels

Problem: the category \mathcal{A} of abelian Polish groups is **not abelian**

Reason: a continuous group homomorphism **need not have closed image**

In 1976 Calvin C. Moore writes about

one final difficulty in considering the cohomology of topological groups which to some extent is incurable, and this is the fact that a continuous group homomorphism need not have closed range.

Table of Contents

- 1 Motivation from topology
- 2 “Completions” of categories of algebraic-topological objects
- 3 Definable refinements of algebraic invariants
 - Finer invariants
 - Richer invariants
 - Rigid invariants

The derived category of abelian Polish groups and its heart

Let \mathcal{A} be the category of abelian Polish groups

The derived category of abelian Polish groups and its heart

Let \mathcal{A} be the category of abelian Polish groups

Let $K(\mathcal{A})$ be the **triangulated category** of complexes over \mathcal{A}

The **derived category** $D(\mathcal{A})$ of \mathcal{A} is obtained from $K(\mathcal{A})$ by “formally inverting” the morphisms whose mapping cone is “strictly exact”

The derived category of abelian Polish groups and its heart

Let \mathcal{A} be the category of abelian Polish groups

Let $K(\mathcal{A})$ be the **triangulated category** of complexes over \mathcal{A}

The **derived category** $D(\mathcal{A})$ of \mathcal{A} is obtained from $K(\mathcal{A})$ by “formally inverting” the morphisms whose mapping cone is “strictly exact”

The **heart** (or **coeur**)

$$\text{LH}(\mathcal{A}) \subseteq D(\mathcal{A})$$

is defined with respect to the canonical (left) t-structure on $D(\mathcal{A})$

The derived category of abelian Polish groups and its heart

Let \mathcal{A} be the category of abelian Polish groups

Let $K(\mathcal{A})$ be the **triangulated category** of complexes over \mathcal{A}

The **derived category** $D(\mathcal{A})$ of \mathcal{A} is obtained from $K(\mathcal{A})$ by “formally inverting” the morphisms whose mapping cone is “strictly exact”

The **heart** (or **coeur**)

$$\text{LH}(\mathcal{A}) \subseteq D(\mathcal{A})$$

is defined with respect to the canonical (left) t-structure on $D(\mathcal{A})$

- $\text{LH}(\mathcal{A})$ is the “smallest” abelian category containing \mathcal{A}
- $D(\mathcal{A}) = D(\text{LH}(\mathcal{A}))$

The derived category of abelian Polish groups and its heart

Let \mathcal{A} be the category of abelian Polish groups

Let $K(\mathcal{A})$ be the **triangulated category** of complexes over \mathcal{A}

The **derived category** $D(\mathcal{A})$ of \mathcal{A} is obtained from $K(\mathcal{A})$ by “formally inverting” the morphisms whose mapping cone is “strictly exact”

The **heart** (or **coeur**)

$$\text{LH}(\mathcal{A}) \subseteq D(\mathcal{A})$$

is defined with respect to the canonical (left) t-structure on $D(\mathcal{A})$

- $\text{LH}(\mathcal{A})$ is the “smallest” abelian category containing \mathcal{A}
- $D(\mathcal{A}) = D(\text{LH}(\mathcal{A}))$

More generally the same applies to any **quasi-abelian** category

An explicit description of the heart of abelian Polish groups

Theorem (L., 2022)

*Explicit description of $\text{LH}(\mathcal{A})$ as a **concrete category***

An explicit description of the heart of abelian Polish groups

Theorem (L., 2022)

Explicit description of $\text{LH}(\mathcal{A})$ as a *concrete category*

Objects: “Formal quotients” G/N of abelian Polish groups by Polish subgroups (the topology of N need not be induced by G)

Morphisms: Group homomorphisms $G/N \rightarrow H/M$ that are *Borel-definable*, i.e. induced by a Borel function $G \rightarrow H$

An explicit description of the heart of abelian Polish groups

Theorem (L., 2022)

Explicit description of $\text{LH}(\mathcal{A})$ as a *concrete category*

Objects: “Formal quotients” G/N of abelian Polish groups by Polish subgroups (the topology of N need not be induced by G)

Morphisms: Group homomorphisms $G/N \rightarrow H/M$ that are *Borel-definable*, i.e. induced by a Borel function $G \rightarrow H$

Techniques: advanced tools and recent results from logic

An explicit description of the heart of abelian Polish groups

Theorem (L., 2022)

Explicit description of $\text{LH}(\mathcal{A})$ as a *concrete category*

Objects: “Formal quotients” G/N of abelian Polish groups by Polish subgroups (the topology of N need not be induced by G)

Morphisms: Group homomorphisms $G/N \rightarrow H/M$ that are *Borel-definable*, i.e. induced by a Borel function $G \rightarrow H$

Techniques: advanced tools and recent results from logic

$\text{LH}(\mathcal{A})$ is the natural framework to develop *definable refinements* of classical homological algebra and algebraic topology

An explicit description of the heart of other categories

Similar descriptions for the heart of other topological-algebraic structures:

- locally compact abelian Polish groups
- totally disconnected locally compact abelian Polish groups
- non-Archimedean abelian Polish groups

An explicit description of the heart of other categories

Similar descriptions for the heart of other topological-algebraic structures:

- locally compact abelian Polish groups
- totally disconnected locally compact abelian Polish groups
- non-Archimedean abelian Polish groups
- R -modules

An explicit description of the heart of other categories

Similar descriptions for the heart of other topological-algebraic structures:

- locally compact abelian Polish groups
- totally disconnected locally compact abelian Polish groups
- non-Archimedean abelian Polish groups
- R -modules
- real/complex Banach spaces \longrightarrow vector spaces with a Banach cover
- Banach spaces over a non-Archimedean valued field
- Fréchet spaces

Table of Contents

- 1 Motivation from topology
- 2 “Completions” of categories of algebraic-topological objects
- 3 Definable refinements of algebraic invariants
 - Finer invariants
 - Richer invariants
 - Rigid invariants

Definable refinements of algebraic invariants

Virtually all group invariants from algebraic topology can be **refined** and seen as invariants taking values in the category of groups with Polish cover

Definable refinements of algebraic invariants

Virtually all group invariants from algebraic topology can be **refined** and seen as invariants taking values in the category of groups with Polish cover

Advantages of the definable versions:

- ① finer invariants (distinguish more spaces, more powerful invariants)
- ② richer invariants (e.g., one can study their **Borel class** and **Borel rank**)
- ③ rigid invariants (fewer automorphisms, better grasp on the **dynamics**)

Table of Contents

- 1 Motivation from topology
- 2 “Completions” of categories of algebraic-topological objects
- 3 Definable refinements of algebraic invariants
 - **Finer invariants**
 - Richer invariants
 - Rigid invariants

Theorem (Bergfalk, L., Panagiotopoulos, 2018–2020)

The following invariants admit *definable refinements*:

- *Steenrod homology of compact spaces*
- *K-homology of compact spaces and of C^* -algebras*
- *Čech cohomology of locally compact spaces*

Theorem (Bergfalk, L., Panagiotopoulos, 2018–2020)

The following invariants admit *definable refinements*:

- Steenrod homology of compact spaces
- \mathbb{K} -homology of compact spaces and of C^* -algebras
- Čech cohomology of locally compact spaces

Furthermore:

- 1 definable Steenrod homology $H_*(-)$ is a complete invariant for solenoids (inverse limits of tori)
- 2 definable \mathbb{K} -homology is a complete invariant for solenoids
- 3 definable Čech cohomology $H^*(-)$ is a complete invariant for mapping telescopes of tori or spheres

Definable homological algebra

The homological invariants

$$\text{Hom}(A, B)$$

$$\text{Ext}(A, B)$$

for countable groups A and B can be seen as groups with a Polish cover.

Definable homological algebra

The homological invariants

$$\text{Hom}(A, B)$$

$$\text{Ext}(A, B)$$

for countable groups A and B can be seen as groups with a Polish cover.

Theorem (Bergfalk, L., Panagiotopoulos, 2019)

The definable homological invariant $\text{Ext}(-, \mathbb{Z})$ is a complete invariant for finite-rank torsion-free abelian groups with no nonzero free summands.

Definable homological algebra

The homological invariants

$$\text{Hom}(A, B)$$

$$\text{Ext}(A, B)$$

for countable groups A and B can be seen as groups with a Polish cover.

Theorem (Bergfalk, L., Panagiotopoulos, 2019)

The definable homological invariant $\text{Ext}(-, \mathbb{Z})$ is a complete invariant for finite-rank torsion-free abelian groups with no nonzero free summands.

Definable $\text{Ext}(-, \mathbb{Z})$ is a fully faithful functor from finite-rank torsion-free abelian groups with no nonzero free summands to groups w/ a Polish cover

Definable homological algebra

The homological invariants

$$\text{Hom}(A, B)$$

$$\text{Ext}(A, B)$$

for countable groups A and B can be seen as groups with a Polish cover.

Theorem (Bergfalk, L., Panagiotopoulos, 2019)

The definable homological invariant $\text{Ext}(-, \mathbb{Z})$ is a complete invariant for finite-rank torsion-free abelian groups with no nonzero free summands.

Definable $\text{Ext}(-, \mathbb{Z})$ is a fully faithful functor from finite-rank torsion-free abelian groups with no nonzero free summands to groups w/ a Polish cover

This does not hold for the purely algebraic Ext .

Spaces with a Banach cover

Theorem

Fix $q < p$ and $q' < p'$

The spaces

$$l_p/l_q \quad \text{and} \quad l_{p'}/l_{q'}$$

are not isomorphic as spaces with a Banach cover when $q \neq q'$.

Spaces with a Banach cover

Theorem

Fix $q < p$ and $q' < p'$

The spaces

$$l_p/l_q \quad \text{and} \quad l_{p'}/l_{q'}$$

are not isomorphic as spaces with a Banach cover when $q \neq q'$.

However, they are always isomorphic as (seminormed) vector spaces.

Table of Contents

- 1 Motivation from topology
- 2 “Completions” of categories of algebraic-topological objects
- 3 Definable refinements of algebraic invariants
 - Finer invariants
 - Richer invariants
 - Rigid invariants

Subobjects

Let $G = \hat{G}/N$ be a group with a Polish cover.

A **subgroup with a Polish cover** H of G is of the form

$$H = \hat{H}/N$$

for some Polishable subgroup \hat{H} of \hat{G} containing N .

Subobjects

Let $G = \hat{G}/N$ be a group with a Polish cover.

A **subgroup with a Polish cover** H of G is of the form

$$H = \hat{H}/N$$

for some Polishable subgroup \hat{H} of \hat{G} containing N .

Such a subgroup H of G has a **Borel class** and a **Borel rank**.

These are by definition the Borel class and the Borel rank of \hat{H} in \hat{G} .

Subobjects

Let $G = \hat{G}/N$ be a group with a Polish cover.

A **subgroup with a Polish cover** H of G is of the form

$$H = \hat{H}/N$$

for some Polishable subgroup \hat{H} of \hat{G} containing N .

Such a subgroup H of G has a **Borel class** and a **Borel rank**.

These are by definition the Borel class and the Borel rank of \hat{H} in \hat{G} .

For example, $\{0\}$ inside \mathbb{R}/\mathbb{Q} is Σ_2^0 and has rank 2

Subobjects

Let $G = \hat{G}/N$ be a group with a Polish cover.

A **subgroup with a Polish cover** H of G is of the form

$$H = \hat{H}/N$$

for some Polishable subgroup \hat{H} of \hat{G} containing N .

Such a subgroup H of G has a **Borel class** and a **Borel rank**.

These are by definition the Borel class and the Borel rank of \hat{H} in \hat{G} .

For example, $\{0\}$ inside \mathbb{R}/\mathbb{Q} is Σ_2^0 and has rank 2

However, $\{0\}$ inside $\mathbb{R}^{\mathbb{N}}/\mathbb{Q}^{\mathbb{N}}$ is Π_3^0 and has rank 3

Solecki subgroups

Theorem (L., 2021, building on Solecki 1999 and Farah–Solecki 2006)

Let G be a group with a Polish cover, and let α be a countable ordinal.

There exists a smallest $\Pi_{1+\alpha+1}^0$ subgroup with a Polish cover $s_\alpha(G)$ of G .

Solecki subgroups

Theorem (L., 2021, building on Solecki 1999 and Farah–Solecki 2006)

Let G be a group with a Polish cover, and let α be a countable ordinal.

There exists a smallest $\Pi_{1+\alpha+1}^0$ subgroup with a Polish cover $s_\alpha(G)$ of G .

Remark

We have that $s_0(G)$ is the closure of $\{0\}$.

Solecki subgroups for Ext of torsion groups

Theorem (L., 2021)

For every countable ordinal α , and torsion groups A and B ,

$$s_\alpha(\text{Ext}(A, B))$$

is equal to the $(1 + \alpha)$ -th Ulm subgroup

$$u_{1+\alpha}(\text{Ext}(A, B))$$

Solecki subgroups for Ext of torsion groups

Theorem (L., 2021)

For every countable ordinal α , and torsion groups A and B ,

$$s_\alpha(\text{Ext}(A, B))$$

is equal to the $(1 + \alpha)$ -th Ulm subgroup

$$u_{1+\alpha}(\text{Ext}(A, B))$$

Corollary (Eilenberg–MacLane, 1942)

The closure of $\{0\}$ in $\text{Ext}(A, B)$ is equal to the first Ulm subgroup, and it is the subgroup corresponding to **pure extensions**.

Solecki subgroups for Ext of torsion groups

Theorem (L., 2021)

For every countable ordinal α , and torsion groups A and B ,

$$s_\alpha(\text{Ext}(A, B))$$

is equal to the $(1 + \alpha)$ -th Ulm subgroup

$$u_{1+\alpha}(\text{Ext}(A, B))$$

Corollary (Eilenberg–MacLane, 1942)

The closure of $\{0\}$ in $\text{Ext}(A, B)$ is equal to the first Ulm subgroup, and it is the subgroup corresponding to **pure extensions**.

Corollary

For torsion groups A, B , $\{0\}$ can have arbitrarily high rank in $\text{Ext}(A, B)$

Solecki subgroups for Ext of torsion groups

Theorem (L., 2021)

For every countable ordinal α , and torsion groups A and B ,

$$s_\alpha(\text{Ext}(A, B))$$

is equal to the $(1 + \alpha)$ -th Ulm subgroup

$$u_{1+\alpha}(\text{Ext}(A, B))$$

Corollary (Eilenberg–MacLane, 1942)

The closure of $\{0\}$ in $\text{Ext}(A, B)$ is equal to the first Ulm subgroup, and it is the subgroup corresponding to **pure extensions**.

Corollary

For torsion groups A, B , $\{0\}$ can have arbitrarily high rank in $\text{Ext}(A, B)$
The problem of classifying extensions can have arbitrarily high complexity.

Solecki subgroups for Ext of torsion-free groups

Theorem (L., 2021)

For a torsion-free groups A, B ,

$$s_1(\text{Ext}(A, B))$$

is the subgroup corresponding to *finite-rank-pure extensions*

Solecki subgroups for Ext of torsion-free groups

Theorem (L., 2021)

For a torsion-free groups A, B ,

$$s_1(\text{Ext}(A, B))$$

is the subgroup corresponding to *finite-rank-pure extensions*

Furthermore, it vanishes when $B = \mathbb{Z}$ or more generally is a subring of \mathbb{Q}

Solecki subgroups for Ext of torsion-free groups

Theorem (L., 2021)

For a torsion-free groups A, B ,

$$s_1(\text{Ext}(A, B))$$

is the subgroup corresponding to *finite-rank-pure extensions*

Furthermore, it vanishes when $B = \mathbb{Z}$ or more generally is a subring of \mathbb{Q}

Corollary

For countable A , *all the finite-rank-pure extensions of A by \mathbb{Z} split.*

Applications to classification by (co)homology

Corollary

If a collection of objects is completely classifiable using as invariants the elements of a Čech cohomology group of a countable CW-complex, then it is also completely classifiable using as invariants countable collections of binary sequences up to tail equivalence

Applications to classification by (co)homology

Corollary

If a collection of objects is completely classifiable using as invariants the elements of a Čech cohomology group of a countable CW-complex, then it is also completely classifiable using as invariants countable collections of binary sequences up to tail equivalence

Thus, **classification by cohomology** is equivalent to reducibility to E_0^ω

Applications to classification by (co)homology

Corollary

If a collection of objects is completely classifiable using as invariants the elements of a Čech cohomology group of a countable CW-complex, then it is also completely classifiable using as invariants countable collections of binary sequences up to tail equivalence

Thus, **classification by cohomology** is equivalent to reducibility to E_0^ω

Corollary (Hopf classification)

Let X be a countable CW-complex with $H^k(X) = 0$ for $k > n$. Then homotopy of maps $X \rightarrow S^n$ is Borel-reducible to E_0^ω .

Applications to the classification problem for extensions

Let X be a compact metrizable space.

We can write $X \approx \lim_n X_n$ where each X_n is a finite polyhedron

Applications to the classification problem for extensions

Let X be a compact metrizable space.

We can write $X \approx \lim_n X_n$ where each X_n is a finite polyhedron

Theorem

Extensions of $C(X)$ by \mathcal{K} are completely classifiable using as invariants countably many binary sequences up to tail equivalence.

Applications to the classification problem for extensions

Let X be a compact metrizable space.

We can write $X \approx \lim_n X_n$ where each X_n is a finite polyhedron

Theorem

Extensions of $C(X)$ by \mathcal{K} are completely classifiable using as invariants countably many binary sequences up to tail equivalence.

Let A be a “well-behaved” C^* -algebra

We can write $A \approx \lim_n A_n$ where A_n has finite-rank \mathcal{K} -groups

Applications to the classification problem for extensions

Let X be a compact metrizable space.

We can write $X \approx \lim_n X_n$ where each X_n is a finite polyhedron

Theorem

Extensions of $C(X)$ by \mathcal{K} are completely classifiable using as invariants countably many binary sequences up to tail equivalence.

Let A be a “well-behaved” C^* -algebra

We can write $A \approx \lim_n A_n$ where A_n has finite-rank K -groups

Theorem

Extensions of A by \mathcal{K} are completely classifiable using as invariants countably many binary sequences up to tail equivalence

Applications to the classification problem for extensions

Let X be a compact metrizable space.

We can write $X \approx \lim_n X_n$ where each X_n is a finite polyhedron

Theorem

Extensions of $C(X)$ by \mathcal{K} are completely classifiable using as invariants countably many binary sequences up to tail equivalence.

Let A be a “well-behaved” C^* -algebra

We can write $A \approx \lim_n A_n$ where A_n has finite-rank K -groups

Theorem

Extensions of A by \mathcal{K} are completely classifiable using as invariants countably many binary sequences up to tail equivalence

This is a **complexity-theoretic consequence** of the UCT for K -homology

Table of Contents

- 1 Motivation from topology
- 2 “Completions” of categories of algebraic-topological objects
- 3 Definable refinements of algebraic invariants
 - Finer invariants
 - Richer invariants
 - Rigid invariants

Rigidity

Groups with a Polish cover are more **rigid** than discrete groups:
they have fewer automorphisms

The reason is that not all group automorphisms are Borel-definable

p -adic numbers

Let \mathbb{Q}_p be the p -adic numbers (seen as additive locally profinite group)

We have a canonical action $\mathbb{Z}[1/p]^\times \curvearrowright \mathbb{Q}_p$ by multiplication

This induces an action $\mathbb{Z}[1/p]^\times \curvearrowright \mathbb{Q}_p/\mathbb{Z}[1/p]$

Ulam stability of p -adics

Theorem (Bergfalk, L., Panagiotopoulos, 2019)

All Borel-definable automorphisms of $\mathbb{Q}_p/\mathbb{Z}[1/p]$ are given by the action

$$\mathbb{Z}[1/p]^\times \curvearrowright \mathbb{Q}_p/\mathbb{Z}[1/p]$$

Thus there exist \aleph_0 Borel-definable automorphisms of $\mathbb{Q}_p/\mathbb{Z}[1/p]$

In contrast, there exist $2^{2^{\aleph_0}}$ automorphisms of $\mathbb{Q}_p/\mathbb{Z}[1/p]$


Solenoids

A solenoid is simply an inverse limit of copies of \mathbb{T}

Solenoids

A solenoid is simply an inverse limit of copies of \mathbb{T}


A concrete geometric realization in \mathbb{R}^3 of a solenoid can be obtained as intersection of a sequence of nested solid tori



Solenoids

A solenoid is simply an inverse limit of copies of \mathbb{T}


A concrete geometric realization in \mathbb{R}^3 of a solenoid can be obtained as intersection of a sequence of nested solid tori



Solenoids

A solenoid is simply an inverse limit of copies of \mathbb{T}


A concrete geometric realization in \mathbb{R}^3 of a solenoid can be obtained as intersection of a sequence of nested solid tori



Solenoids

A solenoid is simply an inverse limit of copies of \mathbb{T}


A concrete geometric realization in \mathbb{R}^3 of a solenoid can be obtained as intersection of a sequence of nested solid tori



Solenoids

A solenoid is simply an inverse limit of copies of \mathbb{T}


A concrete geometric realization in \mathbb{R}^3 of a solenoid can be obtained as intersection of a sequence of nested solid tori



Solenoids

A solenoid is simply an inverse limit of copies of \mathbb{T}

A concrete geometric realization in \mathbb{R}^3 of a solenoid can be obtained as intersection of a sequence of nested solid tori



Solenoid complements

We denote by S^d the one-point compactification of \mathbb{R}^d

Let $X_p \subseteq S^3$ be a geometric realization of the p -adic solenoid

Solenoid complements

We denote by S^d the one-point compactification of \mathbb{R}^d

Let $X_p \subseteq S^3$ be a geometric realization of the p -adic solenoid

Let $[S^3 \setminus X_p, S^2]$ be the space of homotopy classes of maps $S^3 \setminus X_p \rightarrow S^2$

Theorem (Bergfalk, L., Panagiotopoulos, 2020)

There is a Borel-definable bijection

$$[S^3 \setminus X_p, S^2] \cong \mathbb{Q}_p/\mathbb{Z}[1/p]$$

Equivariant classification

Let $\mathcal{E}(S^3 \setminus X_p)$ be the space of **homotopy automorphisms** of $S^3 \setminus X_p$

There is a canonical **Borel-definable action**

$$[S^3 \setminus X_p, S^2] \curvearrowright \mathcal{E}(S^3 \setminus X_p)$$

Equivariant classification

Let $\mathcal{E}(S^3 \setminus X_p)$ be the space of **homotopy automorphisms** of $S^3 \setminus X_p$

There is a canonical **Borel-definable action**

$$[S^3 \setminus X_p, S^2] \curvearrowright \mathcal{E}(S^3 \setminus X_p)$$

Using the **rigidity** of $\mathbb{Q}_p/\mathbb{Z}[1/p]$ we can conclude:

Theorem (Bergfalk, L., Panagiotopoulos, 2020)

The action

$$[S^3 \setminus X_p, S^2] \curvearrowright \mathcal{E}(S^3 \setminus X_p)$$

corresponds to the canonical action

$$\mathbb{Z}[1/p]^\times \curvearrowright \mathbb{Q}_p/\mathbb{Z}[1/p]$$

Equivariant classification

So the problem of classifying the **orbits** of

$$[S^3 \setminus X_p, S^2] \simeq \mathcal{E}(S^3 \setminus X_p)$$

is the same as the problem of classifying the orbits of

$$\mathbb{Z}[1/p]^\times \curvearrowright \mathbb{Q}_p/\mathbb{Z}[1/p]$$

which in turn is the same as the problem of classifying the orbits of

$$\mathbb{Z}[1/p]^\times \ltimes \mathbb{Z}[1/p] \curvearrowright \mathbb{Q}_p$$

Equivariant classification

So the problem of classifying the **orbits** of

$$[S^3 \setminus X_p, S^2] \curvearrowright \mathcal{E}(S^3 \setminus X_p)$$

is the same as the problem of classifying the orbits of

$$\mathbb{Z}[1/p]^\times \curvearrowright \mathbb{Q}_p/\mathbb{Z}[1/p]$$

which in turn is the same as the problem of classifying the orbits of

$$\mathbb{Z}[1/p]^\times \ltimes \mathbb{Z}[1/p] \curvearrowright \mathbb{Q}_p$$

In particular, there exist 2^{\aleph_0} such orbits

Higher dimensions

There are higher-dimensional analogues, where

$$X_p^d \subseteq S^{d+2}$$

is the product of d copies of the p -adic solenoid.

Higher dimensions

There are higher-dimensional analogues, where

$$X_p^d \subseteq S^{d+2}$$

is the product of d copies of the p -adic solenoid.

In this case we have that the Borel-definable action

$$[S^{d+2} \setminus X_p^d, S^{d+1}] \curvearrowright \mathcal{E}(S^{d+2} \setminus X_p^d)$$

corresponds to the action

$$\mathrm{GL}_d(\mathbb{Z}[1/p]) \curvearrowright \mathbb{Q}_p^d / \mathbb{Z}[1/p]^d$$

Measuring the complexity

Using tools from

- **ergodic theory** (superrigidity for profinite actions), and
- **algebraic geometry** (superrigidity for p -adic Lie groups)

one can compare the **Borel complexity** of such actions.

Theorem (Bergfalk, L., Panagiotopoulos, 2019)

The Borel complexity of classifying the orbits of

$$[S^{d+2} \setminus X_p^d, S^{d+1}] \curvearrowright \mathcal{E}(S^{d+2} \setminus X_p^d)$$

or equivalently

$$\mathrm{GL}_d(\mathbb{Z}[1/p]) \ltimes \mathbb{Z}[1/p]^d \curvearrowright \mathbb{Q}_p^d$$

strictly increases with d .

Measuring the complexity

Using tools from

- **ergodic theory** (superrigidity for profinite actions), and
- **algebraic geometry** (superrigidity for p -adic Lie groups)

one can compare the **Borel complexity** of such actions.

Theorem (Bergfalk, L., Panagiotopoulos, 2019)

The Borel complexity of classifying the orbits of

$$[S^{d+2} \setminus X_p^d, S^{d+1}] \curvearrowright \mathcal{E}(S^{d+2} \setminus X_p^d)$$

or equivalently

$$\mathrm{GL}_d(\mathbb{Z}[1/p]) \ltimes \mathbb{Z}[1/p]^d \curvearrowright \mathbb{Q}_p^d$$

strictly increases with d .

*For $d \geq 3$, these problems for different primes are **incomparable** from the perspective of Borel complexity.*

Further directions

Project

*Hierarchies of **phantom maps** corresponding to Solecki subgroups*

Further directions

Project

*Hierarchies of **phantom maps** corresponding to Solecki subgroups*

Project

*Develop definable refinements of algebraic invariants in **coarse geometry***

Further directions

Project

*Hierarchies of **phantom maps** corresponding to Solecki subgroups*

Project

*Develop definable refinements of algebraic invariants in **coarse geometry***

Project

*Definable refinements of group invariants (**bounded cohomology**)*

Further directions

Project

Hierarchies of *phantom maps* corresponding to Solecki subgroups

Project

Develop definable refinements of algebraic invariants in *coarse geometry*

Project

Definable refinements of group invariants (*bounded cohomology*)

Project

Isolate the complexity-theoretic content of the *coarse Baum–Connes conjecture* and of the *Universal Coefficient Theorem* for KK-theory

Further directions

Project

Hierarchies of *phantom maps* corresponding to Solecki subgroups

Project

Develop definable refinements of algebraic invariants in *coarse geometry*

Project

Definable refinements of group invariants (*bounded cohomology*)

Project

Isolate the complexity-theoretic content of the *coarse Baum–Connes conjecture* and of the *Universal Coefficient Theorem* for KK-theory

Project

Construct examples of C^* -algebras and coarse spaces where the UCT and the coarse BC conjecture fail for complexity-theoretic obstructions